

Sanyuania cuneata Zhao & Whatley, 1992
(Ostracoda, Cytheroidea, Cytherideidae) from
Japan: morphology, distribution and
environmental significance

メタデータ	言語: eng 出版者: 公開日: 2021-04-15 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	https://doi.org/10.24517/00061631

This work is licensed under a Creative Commons
Attribution-NonCommercial-ShareAlike 3.0
International License.



MICROPALAEONTOLOGY NOTEBOOK

***Sanyuania cuneata* Zhao & Whatley, 1992 (Ostracoda, Cytheroidea, Cytherideidae) from Japan: morphology, distribution and environmental significance**

GENGO TANAKA

Department of Geology and Mineralogy, Kyoto University Kitashirakawa Oiwakecho, Sakyo-ku, Kyoto 606-8502, Japan

Sanyuania cuneata Zhao & Whatley, 1992 was reported initially from Late Pleistocene core samples from the northern part of the Yellow Sea and Recent sediments collected from supra tidal pools (17 practical salinity unit (psu)) in Xiangshangang Bay, East China Sea (Zhao & Whatley, 1992). Later, specimens were discovered, but not identified, in Lake Nakaumi, Shimane Prefecture, SW Japan (Fig. 1) (as *Cytheridae* gen. et. sp. indet. by Tanaka *et al.* (1998) and *Sanyuania* sp. by Seto *et al.* (1999)). Further examination of these Japanese specimens has clarified that they are conspecific with the Chinese specimens. This has provided an opportunity to review the distribution (Fig. 1) and potential environmental significance of *Sanyuania cuneata* and, for the first time, to describe its appendages (Fig. 2).

Specimens figured herein were recovered from Lake Nakaumi (35°26' 50" N, 133°07'50" E) at a depth of 0.3–6.0 m and salinity of 8–17 psu on 9 September 1998. The surface of the sediment was covered by the byssus of the mussel *Musculista senhousia* (Benson, 1842).

Sanyuania cuneata is probably endemic to the East China Sea and the southwestern part of the Sea of Japan and is a potentially useful palaeoenvironmental indicator of brackish (steno-haline) environments in the area. By using the oxygen isotopic data from planktonic foraminifers and the estimated value of salinity flowing into the Japan Sea, Matsui *et al.* (1998) approximated that the salinity of the surface water of Japan Sea declined to about 20 psu during the Last Glacial Maximum (LGM) and indicated the probability of a land bridge connecting the Asian continent and the Japanese islands at that time. The limited environmental and geographical distribution and the recent fossil record of *S. cuneata* suggest that it took advantage of these shallow, brackish conditions and migrated to the Japanese islands during the LGM. Judging by the fact that




Fig. 2. The chitinous parts of *Sanyuania cuneata*. (1). Antennule. Six articulated segments. (2). Antenna. Four articulated segments. Second segment with long two-segmented exopodite. Fourth segment with two stout clawlike seta. (3). Mandible. Five segmented. Basal segment (coxa, first segment of protopodite) with about six teeth. Second segment of protopodite (basis) with exopodite consisting of two setae, and a plumose seta on posterior distal margin. (4). Maxilla. Branchial plate (exopodite) with 14 plumose setae. Basal podomere bearing palp and three masticatory process. (5). First thoracic leg. (6). Second thoracic leg. (7). Third thoracic leg. (8). Male copulatory organ. Basal capsule rather squared. Hook-shaped short ductus ejaculatorius. 1–8: SUM-CO-1420. Scale = 50 μ m.

the global euryhaline genus *Cyprideis* (Jones) has not been reported in Japan, the Japanese late Cenozoic brackish-water ostracod communities must have connection with the eastern type of Chinese brackish-water ostracod faunas (Zhao & Whatley, 1992).




Fig. 1. Left: SEM photograph of left lateral view of the carapaces of *Sanyuania cuneata* Zhao & Whatley, 1992 (1, female, SUM-CO-1418; 2, male, SUM-CO-1419). Scale = 100 μ m. Right: the distribution of *S. cuneata* in Far East. Fossil (□), Recent (○).

ACKNOWLEDGEMENTS

The author thanks Robin J. Smith (Lake Biwa Museum) for a review of an early draft, Koji Seto (Shimane University) for specimens used in this study. Robin Whatley (University of Wales) and John Gregory (The Natural History Museum) made helpful suggestions.

Manuscript received 3 January 2006

Manuscript accepted 16 June 2006

REFERENCES

Benson, W.H. 1842. Mollusca. In: Cantor, T. (Ed.), General features of Chusan with remarks on the flora and fauna of that island. *Annals and Magazine of Natural History*, 9: 481–493.

- Matsui, H., Tada, R. & Oba, T. 1998. Low-salinity isolation event in the Japan Sea in response to eustatic sea-level drop during LGM: Reconstruction based on salinity-balance model. *The Quaternary Research*, **37**: 221–233.
- Seto, K., Yamaguchi, K. & Tanaka, G. 1999. Capacity of carbonate shell animals for distribution in Lake Nakaumi, Shimane Prefecture, Southwest Japan. *LAGUNA*, **6**: 247–260.
- Tanaka, G., Seto, K. & Takayasu, K. 1998. The relationship between environments and ostracode assemblages from Miho Bay to Lake Shinji. *LAGUNA*, **5**: 81–91.
- Zhao, Q. & Whatley, R. 1992. A revision of the genus *Sanyuania* Zhao and Han, 1980 (Ostracoda, Crustacea) with the description of new species from the Late Cainozoic of China. *Journal of Micropalaeontology*, **11**: 151–158.