

Experimental studies of invasion and metastasis activity of oral squamous cell carcinoma

メタデータ	言語: jpn 出版者: 公開日: 2021-11-12 キーワード (Ja): キーワード (En): 作成者: Yamamoto, Etsuhide メールアドレス: 所属:
URL	https://doi.org/10.24517/00063235

This work is licensed under a Creative Commons
Attribution-NonCommercial-ShareAlike 3.0
International License.



2007 Fiscal Year Final Research Report Summary

Experimental studies of invasion and metastasis activity of oral squamous cell carcinoma

Research Project

Project/Area Number

17390533

Research Category

Grant-in-Aid for Scientific Research (B)

Allocation Type

Single-year Grants

Section

一般

Research Field

Surgical dentistry

Research Institution

Kanazawa University

Principal Investigator

YAMAMOTO Etsuhide Kanazawa University, Graduate School Of Medical Science, Professor (00092445)

Co-Investigator(Kenkyū-buntansha)

NOZAKI Shin-ichi Kanazawa University, Hospital, Assistant professor (10283110)

KAWASHIRI Shuichi Kanazawa University, Graduate School of Medical Science, Instructor (30291371)

Project Period (FY)

2005 - 2007

Keywords

Oral squamous cell carcinoma / Proliferation / Invasion / Metastasis / Adhesion molecule / Matrix / Motility factor / Protease

Research Abstract

We have been studying the mode of cancer invasion as the important prognostic factor in patients with oral squamous cell cancer. Grade 4 of mode of invasion proposed by Jakobsson was subclassified into grades 4C (Cord-like type) and 4D (Diffuse type). The purpose of this study is to immunohistochemically and experimentally elucidate this modified grading system for the mode of invasion. Materials and methods used are human cancer materials, DMBA induced tongue cancer and in vitro and in vivo invasion model by using 3 cell lines; OCS-20 (Gr.3), OSC-19 (Gr. 4C) and HOC-313 (Gr. 4D). In immunostained human cancer materials, the higher the grade of the mode of invasion was, the less continuous the basement membrane was, and the less intercellular adhesion was. Majority of diffuse invasive cancers (Gr. 4C and 4D) co-

expressed membrane type-matrix metalloproteinase and matrix metalloproteinase-2. In cell motility assay, HOC-313 showed the greatest motility (autocrine motility factor; AMF). Cell lines cultured on 3T3 fibroblast-embedded collagen gels showed similar invasion pattern to their origin. Among them, only HOC-313 showed invasion even if fibroblast was not embedded. In DMBA induced tongue cancer, the mode of invasion were observed from Gr.1 to Gr.4C. OSC-20 and OSC-19 implanted orthotopically in the tongue of nude mice grew and metastasized to lymph nodes. Their histologic appearance was similar to that of the origin. HOC-313 grew temporarily for several months by support of matrigel. In recent study, cell selected TSU (Gr.4D) produced lethal tumorigenicity in SCID mouse with similar histology to the origin. This grading system was also proven immunohistochemically and experimentally justifiable.

Research Products (8 results)

	All	2008	2007
	All	Journal Article	Presentation
[Journal Article] Immunohistochemical expressions of E-cadherin and β -catenin crrelate with invasion.			2008 ▾
[Journal Article] Immunohistochemical expressions of E-cadherin and B-catenin crrelate with invasion			2008 ▾
[Journal Article] Copper efflux transpoter contributes to the acquisition of cisplatin resistance in SCC			2007 ▾
[Journal Article] Epigenetic inactivation of IkB kinase in oral carcinomas and tumor progression.			2007 ▾
[Journal Article] Copper efflux transpoter contributes to the acquisition of cisplatin resistance in SCC			2007 ▾
[Journal Article] Epigenetic inactivation of IkB kinase in oral carcinomas and tumor progression			2007 ▾
[Presentation] Behavior of the most invasive oral squamous cell carcinoma(Grade 4D).			2007 ▾
[Presentation] Behavior of the most invasive oral squamous cell carcinoma (Grade 4D)			2007 ▾

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-17390533/173905332007kenkyu_seika_hokoku_

Published: 2010-02-03