Revelation of a extra-cellular matrix and innervations. Of the nerve to the muscle at the time of the muscular tissue repair accompanied by therap eutic exercise

メタデータ	言語: jpn	
	出版者:	
	公開日: 2021-10-15	
	キーワード (Ja):	
	キーワード (En):	
	作成者: Haida, Nobuhide	
	メールアドレス:	
	所属:	
URL	https://doi.org/10.24517/00063809	
This work is licensed under a Creative Commons		

Attribution-NonCommercial-ShareAlike 3.0 International License.



2001 Fiscal Year Final Research Report Summary

Revelation of a extra-cellular matrix and innervations. Of the nerve to the muscle at the time of the muscular tissue repair accompanied by therap eutic exercise

Research Project

Project/Area Number
12832016
Research Category
Grant-in-Aid for Scientific Research (C)
Allocation Type
Single-year Grants
Section
一般
Research Institution
Faculty of Medicine, Kanazawa University
Principal Investigator
HAIDA Nobuhide Kanazawa University, Faculty of Medicine, Professor, 医学部, 教授 (00135089)
Co-Investigator(Kenkyū-buntansha)
YAMAZAKI Toshiaki Kanazawa University, Faculty of Medicine, Assistant, 医学部, 助手 (00220319)
Project Period (FY)
2000 - 2001
Keywords
Muscle repair / Therapeutic exercise / Extracellular matrix / Myoblast / Nerve innervation

Research Abstract

From a functional viewpoint, a repair of the muscular tissue is an important problem on rehabilitation medicine. We examined a role of the therapeutic exercise in the muscular tissue repair. The therapeutic exercise from immediately after a muscle damage induces the activation and proliferation of muscle satellite cell, and earlier appearance of the myotube cell was recognized. And, it became clear that the regeneration of the muscular tissue is accelerated. In the case, fibronectin, laminine and type IV collagen were proved to ambient of the myotube. These extracellular matrix play an important role to the bonding and a shift of the cell. It is conjectured that these are getting involved in a shift and proliferation to a damage site of a myoblast, in order that the constitution of the muscular tissue breaks out in the damage muscle. It is conceivable that even the reinforcement of survival maintenance of the muscle fiber that is not necrosing and resist to the muscle damage to muscle regeneration besides, a control of the proliferation and differentiation of a myoblast is important relation as an adjustment role of muscle regeneration. Not only muscle regeneration is examined only the regeneration but also we need to recognize the interaction between "the neuromusculer system and the extracellular matrix".

Research Products (9 results)

AII O	ther
All Publicat	ions
[Publications] Haida Nobuhide: "Effect of hindlimb re-weighting, low -and high -intensity exercise on disuse atrophy of rat solens muscle"金沢大学医学部保健学科紀要. 24. 1-7 (2000)	~
[Publications] Haida Nobuhide: "Adaptation of slow and fast gastrocnemins muscle to isometric exercise in rat"金沢大学医学部保健学科紀要. 24. 39-44 (2000)	~
[Publications] Yamazaki Toshiaki: "Influence of the time when weight bearing is started on disuse atrophy in rat solens muscle"J. Jpn. Phys. Ther. 4. 13-18 (2001)	~
[Publications] 灰田 信英: "骨格筋の萎縮に対する可逆性の細胞生物学"ボバースジャーナル. 24. 59-62 (2001)	~
[Publications] 灰田 信英: "基礎医学研究の実際"医学書院. 251 (2001)	~
[Publications] Yamazaki T,: "Influence of the time when weight bearing is started on disuse atrophy in rat soleus muscle" Jpn Phys Ther. 4. 13-18 (2001)	~
[Publications] Haida N,: "Adaptation of slow soleus and fast gastrocnemius muscles to isometric exercise in rat"Memoirs Health Sci Med Kanazawa Univ. 24. 39-44 (2000)	~
[Publications] Haida N,: "Cellular aspects on reversibility of skeletal muscle atrophy"Bobath J. 24. 59-62 (2000)	~
[Publications] Haida N,: "Effect of hindlimb re-weighting, low and high-intensity exercise on disuse atrophy of rat soleus Memoirs"Memoirs Health Sci Med Kanazawa Univ. 24. 1-7 (2001)	~

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-12832016/128320162001kenkyu_seika_hokoku_

Published: 2003-09-16