

# 重力異常から検出された“目に見えない断層”の実体の説明

メタデータ	言語: Japanese 出版者: 公開日: 2022-05-30 キーワード: 作成者: メールアドレス: 所属:
URL	<a href="https://doi.org/10.24517/00066120">https://doi.org/10.24517/00066120</a>

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



# 1997 Fiscal Year Final Research Report Summary

---

## Invisible Faults, revealed from Gravity Anomalies.

Research Project

### Project/Area Number

---

08640530

### Research Category

---

Grant-in-Aid for Scientific Research (C)

### Allocation Type

---

Single-year Grants

### Section

---

一般

### Research Field

---

固体地球物理学

### Research Institution

---

Kanazawa University

### Principal Investigator

---

**KONO Yoshiteru** Kanazawa University, Faculty of Science, Professor, 理学部・地球学科, 教授 (80019489)

### Co-Investigator(Kenkyū-buntansha)

---

TAKEUCHI Akira Kanazawa University, Faculty of Science, Professor, 理学部・地球科学科, 教授 (20126494)

### Project Period (FY)

---

1996 – 1997

### Keywords

---

gravity anomaly / horizontal gradient (of gravity anomaly) / invisible faults / Japanese Islands / detection of hidden active faults / basement structure

### Research Abstract

---

Detailed gravity data now cover all over Japan for sufficient enough amount to discuss characteristic spatial variation of gravity anomalies. Distribution of gravity anomalies indicates steep horizontal gradient zones of gravity anomalies. We defined the steep horizontal gradient zone of gravity anomaly (SHGZ of GA) is as of the zone in which the horizontal gradient of 16 mgals or larger sustains for at least 10 km for the first ordered of the horizontal gradient zone.

We analyzed distributions of the SHGZ based on our gravity database by using various image processing techniques. As a result, many SHGZ can be recognized all over Japan, but relatively rare over the Chugoku district. Even though the distribution of the many SHGZ do not coincide with the major geological tectonic boundaries, they appear in parallel within several tens km from them, suggesting that internal relationship between them. Geological and seismological evidences suggest the SHGZ are hidden fault-type geologic structures in a deeper part of the upper crust up to about 10 km deep. Some of them may active faults.

In the Quaternary planes, where are covered by thick soft sediments, there are only limited tools to detect active faults in basement stratum. Gravity anomalies, however have potential to reveal them through detailed gravity measurements.

**URL:** [https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-08640530/086405301997kenkyu\\_seika\\_hokoku\\_](https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-08640530/086405301997kenkyu_seika_hokoku_)

Published: 1999-03-15