

Study on the anti-dementia therapy and quantitative measurement of severity of dementia in the rat model of Alzheimer disease

メタデータ	言語: jpn 出版者: 公開日: 2022-10-24 キーワード (Ja): キーワード (En): 作成者: Ikeda, Kiyonobu メールアドレス: 所属:
URL	https://doi.org/10.24517/00067447

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 International License.



1993 Fiscal Year Final Research Report Summary

Study on the anti-dementia therapy and quantitative measurement of severity of dementia in the rat model of Alzheimer disease

Research Project

Project/Area Number

03670678

Research Category

Grant-in-Aid for General Scientific Research (C)

Allocation Type

Single-year Grants

Research Field

Cerebral neurosurgery

Research Institution

Kanazawa University

Principal Investigator

IKEDA Kiyonobu Department of Neurosurgery, Kanazawa University School of Medicine, 医学部・附属病院, 講師 (80126565)

Project Period (FY)

1991 - 1993

Keywords

Alzheimer disease / nucleus basalis magnocellularis / vagal nodosal ganglion / autotransplantation / acetylcholine marker / event-related potential(P300)

Research Abstract

A unilateral lesion was made in the nucleus basalis magnocellularis (NBM) of the Alzheimer rat model and after one week autotransplantation of the vagal nodosal ganglion (X) was carried out. Four weeks later serial measurements of Acetylcholine (ACh) markers (choline acetyltransferase, CAT ; acetylcholinesterase, AChE ; 3 H-QNB binding activity to muscarine receptor, mAChR) and latency of the P300 (event-related potentials) were carried out for the evaluation of its therapeutic effects. Histological studies by the AChE staining (Tago's method) showed the NBM lesions made by microinjection of the ibotenic acid. Accepted and proliferated viable X tissue was documented in the brain even 4 weeks after X autoplantation. The latency of P300 was 356(SY.+-. ()21.3 msec in normal rats. It was continuously delayd over 400 msec until 4 weeks in the NBM-lesioned rats. Whereas it was found less than 351.0-360.0 msec after 2 weeks in the autotransplanted rats ; indicating significant improvement. The changes of ACh markers was as follows : 1) One weeks after making lesion, CAT activity was reduced to approximately 65% in the cerebral cortex (CC) and 75% in the subcortical structure (SS) on the lesion side compared to that on the contralateral side. From 2 weeks, CAT activity was increased in CC but not in SS. After 3 weeks, AChE activity was reduced to the lowest level and on the contrary mAChR was increased to reach its peak. 2) CAT activity was increased in the CC and SS from 2 weeks after autotransplantation. AChE activity was also increased in SS from 2 weeks and in CC from 3 weeks. However, mAChR was reduced continuously from 2 weeks in the whole brain. From results mentioned above, the conclusions is as follows : 1) In response to the rapidly decreased ACh production caused by

cholinergic deprivation, a kind of compensatory mechanism, such as the increased receptor-binding activity combined with the reduced AChE activity, might take place. 2) Autotransplantation of

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-03670678/036706781993kenkyu_seika_hokoku_

Published: 1995-03-26