

Crystal structures and conformations of 5-benzyl-2-thiohydantoin and its 1-acetylated derivative

メタデータ	言語: eng 出版者: 公開日: 2017-10-03 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/17192

Crystal Structures and Conformations of 5-Benzyl-2-thiohydantoin and Its 1-Acetylated Derivative

Ko-Ki Kunimoto, Masaki Ichtani*, Toshikazu Ogawa, and Soh-ichi Kitoh

Division of Material Sciences, Graduate School of Natural Science and Technology, Kanazawa University, Kakumamachi, Kanazawa, Japan

Akio Kuwae and Kazuhiko Hanai

Graduate School of Natural Sciences, Nagoya City University, Mizuho-ku, Nagoya, Japan

Abstract: The crystal structures of 5-benzyl-2-thiohydantoin (5-BTH) and 1-acetyl-5-benzyl-2-thiohydantoin (1-Ac-5-BTH) have been determined by X-ray diffraction. In the 5-BTH crystals, the enantiomeric (*R*)- and (*S*)-5-BTH molecules are connected to form cyclic dimers *via* the hydrogen bonds of the thioamide and the amide moieties. On the other hand, the intermolecular hydrogen bonds in 1-Ac-5-BTH crystals form an infinite chain. These differences

in the hydrogen bond pattern are also discussed in the IR and Raman spectra. The *ab initio* molecular orbital calculations (Gaussian 03) with 6-31G(d,p) basis set were carried out for 5-BTH and 1-Ac-5-BTH to get the preferred conformation.

Keywords: Crystal structure, molecular conformation, 5-benzyl-2-thiohydantoin, 1-acetyl-5-benzyl-2-thiohydantoin, vibrational spectra, *ab initio* MO calculation

Running title: Crystal Structures of 5-Benzyl-2-thiohydantoin

Address correspondence to Ko-Ki Kunimoto, Division of Material Sciences, Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan. E-mail: kunimoto@sgkit.ge.kanazawa-u.ac.jp

* Present address: Central Research Institute, Ito En, Ltd., 21 Mekami, Makinohara, Shizuoka 421-0516, Japan.

INTRODUCTION

Hydantoins (imidazolidine-2,4-diones), having a 5-membered ring containing a reactive cyclic urea core, form a wide range of biologically active compounds^[1-4]. Their 2-thioxo analogs, 2-thiohydantoins (2-thioxo-imidazolizin-4-one) also display significant biological activities and are employed as established drugs, fungicides or herbicides^[5,6]. Both compounds are considered as useful intermediates in peptide synthesis and structure determination of polypeptides^[7]. Since their biological activities and physicochemical properties are closely related to the electronic structure, conformation and intermolecular interactions, experimental data pertaining to these features are therefore very important. Furthermore, the 2-thiohydantoin furnishes an interesting feature in structural chemistry. This compound carries a thioamide and an amide group in a molecule, which provide equal numbers of proton donor (D) and acceptor (A) in a D-A-D-A sequence. Because of this unique structural feature, 2-thiohydantoins are expected to form intricate

hydrogen bonding networks in crystals. However, there have been few reports on molecular and crystal structures of 2-thiohydantoins compared to hydantoins^[8].

In this work, we have studied the crystal structures and conformations of 5-benzyl-2-thiohydantoin (5-BTH) and 1-acetyl-5-benzyl-2-thiohydantoin (1-Ac-5-BTH). The D-A-D-A hydrogen bonding motif of 5-BTH is modified by acetylation of the NH group in 1-Ac-5-BTH. Differences in IR and Raman spectra between the two compounds were interpreted in terms of changes in the molecular structure and the hydrogen bonding. The theoretical molecular conformations were obtained by *ab initio* MO calculation and compared to the experimental data.

EXPERIMENTAL

Materials

5-BTH was prepared from L-phenylalanine *via* acid hydrolysis of 1-Ac-5-BTH by the thiocyanate method^[9]. 5-BTH and 1-Ac-BTH were obtained as racemic crystals due to racemization of a reaction intermediate^[10]. L-

phenylalanine and ammonium thiocyanate were purchased from Tokyo Kasei Co. Other chemicals were commercial products and used without further purification. L-phenylalanine (5.03 g, 30.0 mmol) was allowed to react with a mixture of ammonium thiocyanate (2.7 g, 35.4 mmol), acetic anhydride (30 ml, 317 mmol) and acetic acid (3.9 ml) at 100 °C for 1 h according to the reported procedure^[9]. A white precipitate of 1-Ac-5-BTH was formed by adding 100 ml of distilled water and subsequent cooling of the solution in a refrigerator and recrystallized from methanol. The 1-Ac-5-BTH obtained was dissolved in 12 M hydrochloric acid and heated at 60 °C for 1 h to yield 5-BTH. The crude product was washed with cold water several times and purified by repeated crystallization from methanol. The purities of these compounds were checked by the elemental analyses and ¹H NMR spectra. 1-Ac-5-BTH: Yield 83%; m.p. 178-180 °C; Found: C, 58.32; H, 4.82; N, 13.59%. Calcd for C₁₂H₁₂N₂O₂S: C, 58.23; H, 4.89; N, 13.59 %. ¹H NMR(270 MHz, DMSO-*d*₆): δ 2.69 (s, 3H, -CH₃), 3.12 (dd, 1H, *J* = 13.5 Hz, 2.7 Hz, -CH₂-), 3.38 (dd, 1H, *J* = 13.5 Hz, 2.7 Hz, -CH₂-), 4.98 (dd, 1H, *J* = 13.5 Hz, 2.7 Hz, -CH-), 6.96-6.99 (m, 2H, -Ph), 7.23-7.31 (m, 3H, -Ph), 12.41 (s, 1H, -NH-). 5-BTH: Yield: 43 %; m.p. 171-174 °C; Found: C, 57.83; H, 4.87; N, 11.12 %. Calcd for C₁₀H₁₀N₂OS: C,

58.05; H, 4.87; N, 11.28 %. ^1H NMR(270 MHz, DMSO- d_6): δ 2.98 (d, 2H, $J = 5.4$ Hz, $-\text{CH}_2-$), 4.56 (t, 1H, $J = 5.4$ Hz, $-\text{CH}-$), 7.15-7.31 (m, 5H, $-\text{Ph}$), 10.06 (s, 1H, $-\text{C}(=\text{S})-\text{NH}-$), 11.43 (s, 1H, $-\text{C}(=\text{O})-\text{NH}-$). The *N*-deuterated 5-BTH and 1-Ac-5-BTH (5-BTH- Nd_2 and 1-Ac-5-BTH- Nd_2) were obtained by exchange reaction with methanol- Od_1 (Merck, 99% atom D).

Spectral Measurements

The IR spectra were recorded on a Perkin Elmer 1650 FT-IR spectrometer as KBr disks, and Nujol and hexachlorobutadiene mulls by averaging 64 scans with a resolution of 4 cm^{-1} . The FT-Raman spectra were obtained on a Perkin-Elmer 2000R spectrometer as powder sealed in a capillary tube. The 1064 nm line of an Elforlight Model L04-2000S Nd:YAG laser was used as the exciting source with an output power of about 200 mW at the sample position. All spectra were accumulated for 60 scans with a resolution of 4 cm^{-1} .

X-ray Crystal Structure Analysis

Single crystals of 5-BTH and 1-Ac-5-BTH suitable for X-ray diffraction analysis were grown by slow evaporation from chloroform and hexane/ethanol solutions, respectively, at room temperature. X-ray diffraction data were obtained on a Rigaku/MSC Mercury CCD diffractometer with a graphite-monochromated Mo K α radiation ($\lambda = 0.7107 \text{ \AA}$). The crystal sample was cooled under a cold nitrogen stream at $-150 \pm 1 \text{ }^\circ\text{C}$ during X-ray exposure to enhance data quality. The data were corrected for both Lorentz and polarization effects. Table 1 summarizes the crystal data and experimental conditions for the crystal structure determination.

The 5-BTH and 1-Ac-5-BTH structures were solved by direct methods using SIR92^[11] and SIR88^[12], respectively. Crystal structure analysis was performed by using the *teXsan* crystallographic software package^[13]. The non-hydrogen atoms were refined anisotropically. All the H-atom positions were found from a difference Fourier map and refined isotropically.ORTEP diagrams were created using the program ORTEP-3^[14].

Quantum Mechanical Calculation

Semiempirical and *ab initio* MO calculations were carried out using the CAChe MOPAC^[15] and the Gaussian 03 set programs^[16], respectively. The starting structures of 5-BTH and 1-Ac-5-BTH were taken from the crystal structure coordinates obtained in this work. The low-energy conformers were searched for the rotation about the C-benzyl bond using the MOPAC AM1 Hamiltonian. In all cases, the PRICISE option was used to provide higher accuracy within this calculation. The low energy conformers were extracted from the MOPAC AM1 calculations and their molecular structures were further optimized by the *ab initio* calculation at the HF level using 6-31G(d,p) basis set.

RESULTS AND DISCUSSION

Crystal Structures of 5-BTH and 1-Ac-5-BTH

The final positional and thermal parameters of 5-BTH and 1-Ac-5-BTH for non-H atoms are presented in Table 2.

Table 3 summarizes the selected structure parameters and the hydrogen bonding geometries. As given in Table 1, 5-

BTH and 1-Ac-5-BTH were crystallized in the monoclinic forms with eight and four molecules, respectively, in a unit cell. The thiohydantoin unit is nearly planar in both molecules, with maximum deviations from planarity of 0.036 and

0.052 Å, respectively, for 5-BTH and 1-Ac5-BTH. As shown in Fig. 1(a), the benzyl moiety of 5-BTH takes the extended conformation with respect to the thiohydantoin ring [C2-C3-C4-C5 -179.8(1) °; N2-C3-C4-C5 66.1(2) °].

The thiohydantoin ring geometries of 5-BTH are comparable to those reported for 2-thiohydantoin and 5,5-diphenyl-2-thiohydantoin^[17-19]. Thus, the C2-O1 and C2-N1 [1.214(2) and 1.374(2) Å] bond distances are in the range observed for the normal *cis*-amide moiety. The C1-S1 bond distance [1.666 (2) Å] is intermediate between those of a C-S bond (1.82 Å) and a C=S bond(1.56 Å), and the C1-N2 distance [1.334(2)Å] is between those of a C-N bond (1.47 Å) and a C=N bond (1.27 Å)^[20].

In contrast to the 5-BTH case, a folded conformation with the aromatic ring over the thiohydantoin ring was found for 1-Ac-5-BTH as shown in Fig. 1(b) [C2-C3-C4-C5 52.2(2) °; N2-C3-C4-C5 -60.7(2) °]. Overall molecular geometries of 1-Ac-5-BTH are similar to those in 1-acetyl-2-thiohydantoin^[21]. The acetyl group on the N2 atom is rotated 11.78(6) ° out of plane from the least square plane of the thiohydantoin ring, and the C11-O2 bond is oriented *trans* to the N2-C1 bond. The bond angle C1-N2-C11 [130.3(1) °] is wider than the C3-N2-C11 [117.7(1) °], resulting

from the repulsion between the S1 atom and the methyl group. The C1-S1[1.645(1) Å] and C1-N2[1.380(2) Å] bond distances of 1-Ac-5-BTH are significantly shorter and longer, respectively, than those of 5-BTH [1.666 (2) and 1.334(2) Å]. Differences in the C1-S1 and C1-N2 bond lengths between 5-BTH and 1-Ac-5-BTH can be rationalized by differences in the resonance structure, as shown in Fig. 2. In the structure of 5-BTH, the zwitterionic canonical forms, Ib and Ic, contribute together with the neutral canonical form Ia to the resonance structure. As a result, the C1-S1 and C1-N2 bonds have single and double bond characters, respectively. On the other hand, in the case of 1-Ac-5-BTH the canonical structures, IIb and IIc, contribute to the resonance structure.

The intermolecular hydrogen bonds which link the molecules are indicated in the molecular packing shown in Figs. 3 and 4. In 5-BTH crystals, the amide and thioamide groups of one molecule form centrosymmetric cyclic dimers with the amide and thioamide groups, respectively, of the adjacent molecules through the intermolecular N-H \cdots O and N-H \cdots S hydrogen bonds [N2 \cdots S1(x-1/2, y+1/2, z), 3.375(2) Å, N2-H2 \cdots S1(x-1/2, y+1/2, z), 178(2) °, N1 \cdots O1(x, y-1, z), 2.831(2)Å, N1-H1 \cdots O1(x, y-1, z), 165(2) °]. The hydrogen bondings form an infinite sheet. On

the other hand, in 1-Ac-5-BTH crystals, the amide N-H of one molecule is hydrogen-bonded to the amide C=O group of another molecule to form an infinite hydrogen bonding chain [N1...O1(-x, y+1/2, -z+1/2), 2.804(2) Å, N1-H1...O1(-x, y+1/2, -z+1/2), 179(2) °]. The S1 atom does not participate in the hydrogen bond system.

IR and Raman Spectra of 5-BTH and 1-Ac-5-BTH and their *N*-Deuterated Analogs

There are some studies on the IR spectra of 2-thiohydantoin derivatives, but no reports have been published on the Raman spectra. Elmore discussed the nature of the thioureide bands^[22], and Poupaert and Bouche studied an IR spectroscopic characterization of these compounds^[23]. Lebedev et al. reported vibrational analyses of 2-thiohydantoin and its 1-acetyl derivative^[24].

Figures 5 and 6 show the IR and Raman spectra of 5-BTH and 1-Ac-5-BTH. The differences in intermolecular hydrogen bondings between 5-BTH and 1-Ac-5-BTH crystals are reflected in their IR and Raman spectra. The typical group frequencies are summarized in Table 4. In the Raman spectrum of 5-BTH, $\nu_{\text{C=O}}$ band is observed at 1724cm^{-1} ,

a lower frequency by 16 cm^{-1} than the corresponding IR band. This frequency difference between the IR and Raman bands is explained in terms of the in-phase and out-of-phase C=O stretchings of the centrosymmetric hydrogen bonding cyclic dimer. Other many IR bands of 5-BTH also have frequencies different from those of the Raman bands, whereas in the 1-Ac-5-BTH spectra the frequency differences are small.

In the IR spectrum of 5-BTH, the C=O band is hardly influenced by *N*-deuteration, but in the Raman spectrum a shift of 18 cm^{-1} to a lower frequency is observed. This shift suggests a possibility that a vibrational coupling with the NH bending within the constituent molecules occurs through the strong hydrogen-bond dimer in the A_g or B_g crystal modes. 5-BTH crystals belong to a space group C_{2h}^6 ($C2/c$); the A_g and B_g crystal modes are Raman-active and the A_u and B_u crystal modes are IR-active. For 1-Ac-5-BTH, the ring $\nu\text{C=O}$ is observed as a doublet (1756 and 1730 cm^{-1}) in the Raman spectrum. This splitting is probably caused by the crystal field; since a space group is C_{2h}^5 ($P2_1/c$), the two crystal modes (A_g and B_g) appear in the Raman spectrum. Although no clear splitting is observed in this region of the IR spectrum, a weak shoulder band occurs at 1735 cm^{-1} a lower frequency side of the very strong 1748 cm^{-1} band,

suggesting the presence of this splitting. In fact, a splitting clearly appears at 1747 and 1720 cm^{-1} in the IR spectrum of 1-Ac-5-BTH- Nd_1 , corresponding to the A_u and B_u crystal modes. A lower component of the doublet is shifted by about 15 cm^{-1} in the both spectra on N -deuteration. This behavior could be attributed to a difference in coupling between the crystal modes.

The $\nu\text{C=O}$ of the acetyl group of 1-Ac-5-BTH is ascribed to the 1705 cm^{-1} band in the IR spectrum and to the 1704 cm^{-1} band in the Raman spectrum. Although the bond length of the acetyl C=O is slightly shorter than that of the ring C=O (Table 3), this assignment is reasonable, since this band is not shifted at all by N -deuteration. Contrarily, for 1-acetyl-2-thiohydantoin Lebedev et al. assigned the bands at 1788 cm^{-1} and 1714 cm^{-1} to the acetyl C=O and the ring C=O band, respectively, stating that the presence of the acetyl group leads to a decrease in the ring $\nu\text{C=O}$ (from 1738 cm^{-1} in thiohydantoin to 1714 cm^{-1} in the acetyl derivative)^[24b]. However, their assignment should be revised, because the following effects are expected; in such five-membered ring carbonyl compounds C=O stretching frequencies increase owing to ring strain, and

furthermore in 1-acetyl-2-thiohydantoin the acetyl group attracts electrons from the ring, so that the ring C=O frequency is higher than that of the acetyl C=O.

Quantum Mechanical Calculation

The molecular conformations of 5-BTH and 1-Ac-5-BTH were searched for the rotation about the C3-C4 bond using the MOPAC AM1 method. The torsion angle (C2-C3-C4-C5) was varied between -180° and 180° . As a result, three local minima were found for 5-BTH and 1-Ac-5-BTH which correspond to the I-A, I-B and I-C conformers for 5-BTH and the II-A, II-B and II-C for 1-Ac-5-BTH. The molecular structures of these low energy conformers were further optimized by the *ab initio* calculation at the HF level using 6-31G(d,p) basis set. Table 5 gives selected structural geometries and total energies of these conformers calculated by the *ab initio* MO, together with the X-ray data. The I-B and II-A forms were obtained as the lowest energy conformers for 5-BTH and 1-Ac-5-BTH, respectively, although the energy difference is very small (1.33-2.90 kcal/mol for 5-BTH and 1.86-4.33 kcal/mol for 1-Ac-5-BTH). The I-A, I-B and I-C forms and the II-A, II-B and II-C have similar bond lengths and bond angles, respectively, but the torsion

angles are very different. The I-B form of 5-BTH has an extended conformation with the N2-C3-C4-C5 torsion angle of 65.02° , whereas the II-A form of 1-Ac-5-BTH takes a folded conformation with the N2-C3-C4-C5 torsion angle of -55.92° . These values are close to those obtained by the X-ray analysis [$66.1(2)^\circ$ and $-60.7(2)^\circ$]. These observations indicate that these molecules take rather relaxed conformations in the crystal phase. For 5-BTH, the calculated C2-O1 (C=O) and C1-N1 bond lengths in the I-B form are shorter than those in the X-ray structure [1.186 and 1.370 Å compared to 1.214 and 1.386 Å, respectively]. 1-Ac-5-BTH in the II-A form also shows similar features [1.186 and 1.361 Å compared to 1.217 and 1.380 Å, respectively]. These differences in the C2-O1 and C1-N1 bond lengths between the experimental and the calculated values can be rationalized by the NH...O intermolecular hydrogen bonds formed both in 5-BTH and 1-Ac-5-BTH crystals. These intermolecular hydrogen bonds in the crystals lead to the electronic redistribution of the 2-thiohydantoin ring, so that the C2-O1 and C1-N1 bond distances become longer than those obtained by the quantum chemical calculations for the isolated molecules. In either compound there is no significant difference in the C=S bond length.

ACKNOWLEDGEMENTS

The authors are grateful to Mrs. M. Arai of Kanazawa University for the NMR measurements and to Mrs. M.

Hosokawa and Mrs. M. Hayashi of Gifu Pharmaceutical University for the elemental analyses.

REFERENCES

1. Knabe, J.; Baldauf, J.; Ahlhelm, A. Racemates and enantiomers of basic substituted 5-phenylhydantoins. Syntheses and antiarrhythmic activity. *Pharmazie* **1997**, *52*, 912-919.
2. Scholl, S.; Koch, A.; Henning, D.; Kempfer, G.; Kleinpeter, E. The influence of structure and lipophilicity of hydantoin derivatives on anticonvulsant activity. *Struct. Chem.* **1999**, *10*, 355-366.
3. Rodgers, T. R.; LaMontagne, M. P.; Markovac, A.; Ash, A. B. Hydantoins as antitumor agents. *J. Med. Chem.* **1977**, *20*, 591-594.

4. Bac, P.; Maurois, P.; Dupont, C.; Pages, N.; Stables, J. P.; Gressens, P.; Evrard, P.; Vamecq, J. Magnesium deficiency-dependent audiogenic seizures (MDDASs) in adult mice: a nutritional model for discriminatory screening of anticonvulsant drugs and original assessment of neuroprotection properties. *J. Neurosci.* **1998**, *18*, 4363-4373.
5. Ghoneim, K. M.; El-Telbany, F.; Ismail, M. A. Hydantoin and thiohydantoin derivatives as potential antimicrobial agents: Part I. *Egypt. J. Pharm. Sci.* **1987**, *28*, 77-86.
6. Marton, J.; Enisz, J.; Hosztafi, S.; Timar, T. Preparation and fungicidal activity of 5-substituted hydantoins and their 2-thio analogs. *J. Agric. Food Chem.* **1993**, *41*, 148-152.
7. Bodanszky, M. *Principles of Peptide Synthesis*; Springer Verlag: Berlin, 1993; 95-.
8. Yu, F.-L.; Schwalbe, C. H.; Watkin, D. J. Hydantoin and hydrogen-bonding patterns in hydantoin derivatives. *Acta Cryst.*, **2004**, *C60*, o714-o717.

9. Schlack, P.; Kumpf, W. A new method for ascertaining the constitution of peptides. *Z. physiol. Chem.* **1926**, *154*, 125-170.
10. Bailey, J. M.; Shively, J. E. Carboxy-terminal sequencing: formation and hydrolysis of C-terminal peptidylthiohydantoins. *Biochemistry* **1990**, *29*, 3145-3156.
11. Altomare, A.; Cascarano, G.; Giacovazzo, G.; Guagliardi, A.; Burla, M.C.; Polidori, G.; Camalli, M. SIR92 - a program for automatic solution of crystal structures by direct methods. *J. Appl. Cryst.* **1994**, *27*, 435-435.
12. Burla, M. C.; Camalli, M.; Cascarano, G.; Giacovazzo, C.; Polidori, G.; Spagna, R.; Viterbo, D. SIR88 - a direct-methods program for the automatic solution of crystal structures. *J. Appl. Crystallogr.* **1989**, *22*, 389-393.
13. teXsan, Crystal Structure Analysis Package, Molecular Structure Corp., The Woodlands, Texas. 1985.
14. Farrugia, L. J. ORTEP-3 for windows - a version of ORTEP-III with a graphical user interface (GUI). *J. Appl. Crystallogr.*, **1997**, *30*, 565.

15. Quantum CAChe 5.0 for Windows, Fujitsu Ltd., Japan, 2002.
16. Gaussian 03, Revision A.1: Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, J. A., Jr.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; Pople, J. A.; Gaussian, Inc., Pittsburgh PA, 2003.

17. Walker, L. A.; Folting, K.; Merritt, L. L., Jr. Crystal structure of 2-thiohydantoin. *Acta Crystallogr.*, **1969**, *B25*, 88-93.
18. Devillanova, F. A.; Isaia, F.; Verani, G.; Battaglia, L. P.; Corradi, A. B. HOSE calculations and structural analyses of some 5,5-dimethylimidazolidine-2,4-dichalcogen derivatives. *J. Chem. Res.*, **1987**, *6*, 192-193.
19. Roszak, A. W.; Weaver, D. F. 5,5-Diphenyl-2-thiohydantoin. *Acta Crystallogr.* **1998**, *C54*, 1168-1170.
20. Sutton, L. E. *Tables of Interatomic Distances and Configurations in Molecules and Ions: Supplement*; The Chemical Society: London, 1965; 1956-1959.
21. Casas, J. S.; Castineiras, A.; Couce, D.; Playa, N.; Sordo, J.; Verela, J. M. 1-acetyl-2-thiohydantoin. *Acta Cryst.* **1998**, *C54*, 427-428.
22. Elmore, D.T. Infrared spectra of some 2-thiohydantoin : the nature of the thioureide band. *J. Chem. Soc.* **1958**, 3489-3492.

23. Poupaert, J.; Bouche, R. J. IR spectroscopic characterization of 2-thiohydantoins and 2-thiobarbiturates. *J. Pharm. Sci.* **1976**, *65*, 1258-1260.
24. (a) Lebedev, R.S.; Chumakova, R.P.; Yukhimets, V.N.; Yakimenko, V.I. Calculation and study of the infrared absorption spectrum of thiohydantoin. *Izvestiya Vysshikh Uchebnykh Zavedenii Fizika* **1970**, *13*, 29-33. (b) Lebedev, R.S.; Chumakova, R.P.; Yukhimets, V.N.; Yakimenko, V.I. Infrared absorption spectra of N-acetyl-2-thiohydantoin. *Izvestiya Vysshikh Uchebnykh Zavedenii Fizika* **1971**, *14*, 109-114.

Table 1. Crystal data and structure refinement

Compound	5-BTH	1-Ac-5-BTH
Color / shape	colorless/prism	colorless/prism
Chemical formula	C ₁₀ H ₁₀ N ₂ OS	C ₁₂ H ₁₂ N ₂ O ₂ S
Formula weight	206.26	248.30
Temperature, K	123	123
Crystal system	monoclinic	monoclinic
Space group	<i>C2/c</i> (<i>C</i> _{2h} ⁶)	<i>P2₁/c</i> (<i>C</i> _{2h} ⁵)
Unit cell dimensions	<i>a</i> =13.368(3)Å <i>b</i> =5.7553(8)Å <i>c</i> =25.882(5)Å <i>β</i> =96.1840(9)°	<i>a</i> =11.508(2)Å <i>b</i> =13.396(3)Å <i>c</i> =7.714(2)Å <i>β</i> =95.380(4)°
Volume, Å ³	1979.6(6)	1184.1(4)
<i>Z</i>	8	4
Density (calculated), Mg/m ³	1.384	1.393
Absorption coefficient, mm ⁻¹ mm ⁻¹	0.293	0.264
Diffractometer	Rigaku/ MSC Mercury	Rigaku/ MSC Mercury
<i>θ</i> range for data collection, deg	4.8-55.0	15.4-55.0
Reflections measured	7262	12329
Independent reflections	2052 (<i>R</i> _{int} =0.043)	2742 (<i>R</i> _{int} =0.034)
Observed reflections	1700 [<i>I</i> >1.20σ(<i>I</i>)]	2270 [<i>I</i> >1.20σ(<i>I</i>)]
Data/ restraints/ parameters	1700 / 0 / 167	2270 / 0 / 202
Goodness of fit	1.46	1.21
Final <i>R</i> indices [<i>I</i> >1.20σ(<i>I</i>)]	<i>R</i> =0.040, <i>wR</i> = 0.051	<i>R</i> =0.035, <i>wR</i> = 0.044
Largest diff. peak and hole, e/Å ³	0.34 -0.16	0.26 -0.14

Table 2. Fractional atomic coordinates and equivalent isotropic thermal parameters for non-hydrogen atoms

Atom	X	Y	Z	$B_{\text{eq}}/\text{\AA}^2$
5-BTH				
S1	0.13862(3)	-0.08450(7)	-0.05987(1)	1.961(10)
O1	0.05475(9)	0.5058(2)	0.07014(4)	2.24(2)
N1	0.08229(9)	0.2576(2)	0.00265(5)	1.77(2)
N2	0.1906(1)	0.0132(3)	0.04177(5)	2.23(3)
C1	0.1381(1)	0.0588(3)	-0.00386(6)	1.76(3)
C2	0.0951(1)	0.3381(3)	0.05290(6)	1.94(3)
C3	0.1734(1)	0.1803(3)	0.08240(6)	1.98(3)
C4	0.1385(1)	0.0660(3)	0.13036(6)	2.03(3)
C5	0.2198(1)	-0.0878(3)	0.15807(5)	1.98(3)
C6	0.1917(1)	-0.2970(3)	0.17868(6)	2.30(3)
C7	0.2622(2)	-0.4430(3)	0.20537(6)	2.66(3)
C8	0.3626(1)	-0.3810(3)	0.21165(7)	2.69(8)
C9	0.3930(1)	-0.1743(4)	0.19102(7)	2.85(4)
C10	0.3213(1)	-0.0264(3)	0.16435(6)	2.48(3)
1-Ac-5-BTH				
S1	0.84753(3)	0.03491(3)	0.12011(4)	1.454(8)
O1	0.88337(10)	-0.25409(7)	-0.2655(1)	1.54(2)
O2	0.8813(1)	0.11039(8)	-0.4531(1)	2.04(2)
N1	0.8638(1)	-0.12893(9)	-0.0683(2)	1.13(2)
N2	0.8535(1)	0.00571(8)	-0.2336(1)	1.11(2)
C1	0.8542(1)	-0.02630(10)	-0.0636(2)	1.06(2)
C2	0.8721(1)	-0.16623(10)	-0.2307(2)	1.12(3)
C3	0.8589(1)	-0.07901(10)	-0.3551(2)	1.14(3)
C4	0.7470(1)	-0.0898(1)	0.4802(2)	1.45(3)
C5	0.6393(1)	-0.1080(1)	-0.3873(2)	1.54(3)
C6	0.5963(2)	-0.2041(1)	-0.3703(3)	2.58(4)
C7	0.4970(2)	-0.2206(2)	-0.2839(3)	3.71(5)
C8	0.4404(2)	-0.1424(2)	-0.2135(3)	3.14(4)
C9	0.4820(1)	-0.0462(2)	-0.2286(2)	2.51(4)
C10	0.5806(1)	-0.0293(1)	-0.3155(2)	1.94(3)
C11	0.8601(1)	0.1030(1)	-0.3020(2)	1.42(3)
C12	0.8408(2)	0.1914(1)	-0.1900(2)	1.84(3)

$$B_{eq} = (8/3) \pi^2 (U_{11}(aa^*)^2 + U_{22}(bb^*)^2 + U_{33}(cc^*)^2 + 2U_{12}aa^*bb^*\cos \gamma + 2U_{13}aa^*cc^*\cos \beta + 2U_{23}bb^*cc^*\cos \alpha)$$

Table 3. Selected bond lengths(Å), and bond and torsion angles(°) for 5-BTH and 1-Ac-5-BTH

	5-BTH	1-Ac-5-BTH
<i>Bond lengths</i>		
S1-C1	1.666(1)	1.645(2)
O1-C2	1.214(2)	1.217(2)
N1-C1	1.386(2)	1.380(2)
N1-C2	1.374(2)	1.360(2)
N2-C1	1.334(2)	1.380(2)
N2-C3	1.462(2)	1.477(2)
C2-C3	1.527(2)	1.510(2)
N2-C11		1.415(2)
O2-C11		1.214(2)
C11-C12		1.494(2)
<i>Bond angles</i>		
C1-N1-C2	112.3(1)	113.8(1)
C1-N2-C3	113.5(1)	111.6(1)
S1-C1-N1	124.0(1)	122.0(1)
S1-C1-N2	128.9(1)	131.9(1)
N1-C1-N2	107.1(1)	106.1(1)
O1-C2-N1	126.9(1)	125.2(1)
O1-C2-C3	126.7(1)	127.9(1)
N1-C2-C3	106.4(1)	106.8(1)
N2-C3-C2	100.6(1)	101.5(1)
N2-C3-C4	112.8(1)	112.7(1)
C2-C3-C4	114.3(1)	110.7(1)
C3-C4-C5	111.7(1)	113.2(1)
O2-C11-N2		117.0(1)
N2-C11-C12		119.7(1)
O2-C11-C12		123.2(1)
<i>Torsion angles</i>		
S1-C1-N1-C2	178.5(1)	-178.0(1)
S1-C1-N2-C3	179.2(1)	-178.8(1)
O1-C2-N1-C1	-178.7(2)	178.6(1)
O1-C2-C3-N2	179.0(2)	-178.0(1)

O1-C2-C3-C4	57.9(2)	62.1(2)
N1-C1-N2-C3	-0.1(2)	1.9(1)
N1-C2-C3-N2	-3.1(2)	4.6(1)
N1-C2-C3-C4	-124.2(1)	-115.2(1)
N2-C1-N1-C2	-2.2(2)	1.4(2)
N2-C3-C4-C5	66.1(2)	-60.7(2)
C1-N1-C2-C3	3.4(2)	-4.0(2)
C1-N2-C3-C2	1.9(2)	-4.0(1)
C1-N2-C3-C4	124.1(1)	114.4(1)
C2-C3-C4-C5	-179.8(1)	52.2(2)
O2-C11-N2-C1		165.9(1)
C3-N2-C11-C12		172.3(1)

D-H...A	D-H (Å)	H...A(Å)	D...A(Å)	D-H...A(°)
5-BTH				
N2-H...S1 ⁱ⁾	0.80(2)	2.58(2)	3.375(2)	179(2)
N1-H...O1 ⁱⁱ⁾	0.78(2)	2.07(2)	2.831(2)	165(2)
1-Ac-5-BTH				
N1-H1...O1 ⁱⁱⁱ⁾	0.80(2)	2.00(2)	2.804(2)	179(2)

Symmetry codes: (i) $x-1/2, y+1/2, z$ (ii) $x-1/2, y-1, z$ (iii) $-x, y+1/2, -z+1/2$

Estimated standard deviations in the least significant figure are given in parentheses.

Table 4. Some characteristic frequencies of 5-BTH and 1-Ac-5-BTH and their *N*-deuterated compounds

5-BTH		5-BTH- Nd_2		1-Ac-5-BTH		1-Ac-5-BTH- Nd_1		Assignment
IR (KBr disk)	Raman	IR (Nujol)	Raman	IR (KBr disk)	Raman	IR (Nujol)	Raman	
3175 vs	3163 w	2365 s	2376 w					N-H(N-D), thioamide
3102 s	--	--	--	3106 m	3101 vw	2318 m	2319 vw	N-H(N-D), amide
1740 vs	1724 w	1738 vs	1706 m	1747 vs	1756mw	1746 m	1754 mw	C=O, amide
				1735 sh	1731 mw	1720 w	1717 ms	(C=O, amide, doublet)
				1704 vs	1704 m	1705 vs	1705 w	C=O, acetyl
1549 s	1522 vw	1496 m	--	1464 s	1462 w	1419 s	1422 vw	C-N+ N-H or C-N (thioureide band)

Table 5. Comparison of structural parameters and total energies obtained by *ab initio* MO calculation and X-ray analysis

Parameters	5-BTH				1-Ac-5-BTH			
	Calculated			Experimental	Calculated			Experimental
	I-A	I-B	I-C		II-A	II-B	II-C	
Bond length(Å)								
S1-C1	1.656	1.656	1.656	1.666(1)	1.651	1.653	1.652	1.645(2)
O1-C2	1.186	1.186	1.186	1.214(2)	1.186	1.186	1.186	1.217(2)
N1-C1	1.369	1.370	1.367	1.386(2)	1.361	1.362	1.359	1.380(2)
N1-C2	1.370	1.373	1.375	1.374(2)	1.370	1.374	1.376	1.360(2)
N2-C1	1.334	1.333	1.334	1.334(2)	1.369	1.365	1.369	1.380(2)
N2-C3	1.448	1.448	1.451	1.462(2)	1.474	1.472	1.476	1.477(2)
C2-C3	1.523	1.523	1.527	1.527(2)	1.511	1.512	1.512	1.510(2)
Torsion angles(°)								
N2-C3-C4-C5	-56.94	65.02	165.60	66.1(2)	-55.92	67.75	177.33	-60.7(2)
C2-C3-C4-C5	58.35	179.04	-80.69	-179.8(1)	65.29	175.72	-84.00	52.2(2)
Total energy(a.u.)	-	-	-		-	-	-	
	965.84359	965.84571	965.84107		1117.62013	1117.61323	1117.61717	
Energy difference ΔE (kcal/mol)	1.33	0.00	2.90		0.00	4.33	1.86	

Figure captions

Figure 1. ORTEP drawings of 5-BTH (a) and 1-Ac-BTH (b). The *R*-antipode of a racemic pair is shown with the atom numbering.


Figure 2. Resonance structures related to the thioamide group in BTH (Ia, Ib, Ic) and 1-Ac-5-BTH (IIa, IIb, IIc).

Figure 3. Perspective views of intermolecular hydrogen bonds in BTH crystals..

Figure 4. Perspective views of intermolecular hydrogen bonds in 1-Ac-BTH crystals.

Figure 5. IR (a) and Raman (b) spectra of 5-BTH.

Figure 6. IR (a) and Raman (b) spectra of 1-Ac-5-BTH.



(Fig. 1)




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6