

Influence of phosphate and iron ions in selective uptake of arsenic species by water fern (*Salvinia natans* L.)

メタデータ	言語: eng 出版者: 公開日: 2017-10-03 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/13361

Influence of Phosphate and Iron Ions in Selective Uptake of Arsenic Species by Water Fern (*Salvinia natans* L.)

M. Azizur Rahman^{*1}; H. Hasegawa^{*1}; K. Ueda¹; T. Maki¹; M. Mahfuzur Rahman²

¹Graduate School of Natural Science & Technology, Kanazawa University, Kakuma, Kanazawa

920-1192, Japan; ²Department of Botany, Faculty of Biological Sciences, Jahangirnagar

University, Savar, Dhaka-1342, Bangladesh.

***Corresponding author**

E-mail: hhiroshi@t.kanazawa-u.ac.jp (H. Hasegawa)

arahman@stu.kanazawa-u.ac.jp (M. A. Rahman)

Tel/Fax: 81-76-234-4792

Abstract:

In the present study, the effect of phosphate ion and iron hydroxides (Fe-plaques) on the selective uptake of arsenic species by water fern (*Salvinia natans* L.) was investigated. The plants were grown for 5 days in aqueous Murashige and Skoog (MS) culture media modified in arsenic and phosphate concentrations. Arsenic accumulations in *Salvinia natans* L. increased with the increase of arsenate and DMAA concentrations in the culture solutions. Compared to the control treatment, *Salvinia natans* L. accumulated significantly higher amount of arsenic from phosphate deficient solutions, when the source was arsenate. However, arsenic uptake was not affected significantly by phosphate, when the source was dimethylarsinic acid (DMAA). From solutions modified in 100 μM of phosphate and 4.0 μM of either arsenate or DMAA, the *Salvinia natans* L. accumulated 0.14 ± 0.02 and $0.02 \pm 0.00 \mu\text{mol (g dry weight)}^{-1}$ of arsenic, respectively. In contrast, plants accumulated 0.24 ± 0.06 and $0.03 \pm 0.00 \mu\text{mol (g dry weight)}^{-1}$ of arsenic from solutions containing 4.0 μM of either arsenate or DMAA in the absence of phosphate, respectively. Thus, it is reasonable to state that increasing phosphate concentration in culture solutions decreases the arsenic uptake into the water fern significantly, when the source was arsenate. Moreover, arsenic and phosphate content in plant tissue correlated significantly ($r = -0.66$; $p < 0.05$), when initial source was arsenate while there were no correlation between arsenic and phosphate, when initial source was DMAA ($r = -0.077$; $p > 0.05$). Similarly, significant correlation was observed between arsenic and iron content in plant tissues ($r = 0.66$; $p < 0.05$), when initial source was arsenate while the correlation was not significant ($r = 0.23$; $p < 0.05$), when initial source was DMAA. The results indicate the adsorption of arsenate on Fe-plaques of aquatic plant surfaces. Further, the study demonstrates that the DMAA uptake mechanisms into the water fern are different from those of arsenate.

Keywords: Arsenate; DMAA; Uptake; Physico-chemical Adsorption; Water Fern (*Salvinia natans* L.); Phosphate; Phytofiltration.

Introduction:

Arsenic is one of the toxic environmental pollutants which have recently attracted attention because of its chronic and epidemic effects to the human health through widespread water and crop contamination. Natural release of arsenic from aquifer rocks has been reported in Bangladesh [1-4], West Bengal, India [5, 6]. Geogenic contamination of arsenic in aquifer rocks has also been reported in Thailand [7], Vietnam, inner Mongolia, Greece, Hungary, USA, Ghana, Chile, Argentina and Mexico [8, 9]. Beside the large-scale arsenic pollution in soils, water pollution by geogenic arsenic has been a great health problem in many countries [2, 4, 6].

Phytoremediation, a plant based green technology, becomes promising to remediate the environmental pollution due to some unavoidable limitations of traditional technologies. It is relatively inexpensive, eco-friendly and proven effective in few cases [10]. Although the arsenic uptake into the plants occurs primarily through the root system, it is not readily translocated to the shoots and the edible parts of all plants. Few terrestrial plant species, such as *Agrostis castellana*, *Agrostis delicatula* [11], *Bidens cynapiifolia* [12], Chinese brake fern (*Pteris vittata* L.) [13] and silver fern (*Pityrogramma calomelanos* L.) [14] accumulate high concentration of arsenic in their shoots and edible parts even though the background concentration in soil is low [13]. In particular, Chinese brake fern removes a significant amount of arsenic from soil [14,15], and stores in the fronds [14, 16]. Arsenic accumulation in aquatic plants, such as *Spirodella polyrhiza* L. [5], *Lemna gibba* L. [17, 18], *Hydrilla verticillata* [19], *Lepidium sativum* [20] has also been reported in literatures.

Arsenate; As (V) and arsenite; As (III) are the inorganic forms in the oxic aquatic systems. Arsenate predominates and arsenite is oxidized to arsenate in the oxic aquatic systems [21]. The use of aquatic macrophytes or other floating plants in phytoremediation technology is commonly known as phytoextraction. This clean up process involves biosorption and accumulation of

pollutants. Recently, aquatic macrophytes and some other small floating plants have been investigated for the remediation of wastewater contaminated with Cu, Cd(II) and Hg(II) [22, 23, 24]. The encouraging results of metal uptake capacity by aquatic plants [22-28] gained the attention of researchers and scientists to use them in phytoremediation technology.

Water fern (*Salvinia natans* L.) is a free floating freshwater macrophyte, which grows rapidly in ponds, lakes, ditches, and wastewater bodies mostly in southern Asian countries affected by arsenic especially in Bangladesh, West Bengal, India. Previously, the *Salvinia natans* L. was tested for Hg (II) [24] and Cu (II) [28] removal. In the present study, the authors investigated the effect of phosphate concentrations on arsenate and DMAA uptake and biosorption by *Salvinia natans* L. from aqueous culture solution. The arsenate was selected because it is the predominant inorganic species in oxic aquatic systems [21]. An organic species (DMAA) was also selected to compare the response of the plant to both organic (DMAA) and inorganic (arsenate) species uptake and biosorption in the plant.

Materials and Methods:

Plant Cultivation

The *Salvinia natans* L. were collected from rice field of Manikgonj of Dhaka, Bangladesh and stock-cultured in a green house for two weeks. The experiment was conducted in an incubator for a 5 days period with the conditions being set as 14/10 h light/dark schedule, 100-125 $\mu\text{E m}^{-2} \text{s}^{-1}$ light intensity, 75% humidity, 22 and 20 (± 2) °C temperatures for day and night, respectively.

Plants in the incubator were grown on modified murashige and skoog (MS) culture media where modifications were in phosphorus and arsenic concentrations (Table 1). The modified culture

solutions had either 50 or 100 μM of PO_4^{3-} . Either arsenate or DMAA were added to the modified solutions at the rate of 1.0, 2.0 and 4.0 μM prepared from $\text{Na}_2\text{HAsO}_4 \cdot 7\text{H}_2\text{O}$ and $(\text{CH}_3)_2\text{AsO}_2\text{Na} \cdot 3\text{H}_2\text{O}$, respectively. The control solution contains neither arsenic nor PO_4^{3-} .

Inoculation Procedure

Before inoculation, *Salvinia natans* L. strains from stock-culture were washed three times with DI water. 200-ml polystyrene test vessels (118 X 86 X 60 mm) were used for the experiments. About 10 individual plants were inoculated in each of 200-ml test vessels containing 100 ml of test solution. The pH during the experiments was maintained at 5.5 through adjustment with the addition of either 0.1 M HCl or 0.1 M NaOH. Changes in volume of culture solutions during the experiment from evaporation and accumulation were compensated by adding DI water equivalent to the volume difference in every 2 days throughout the experiment.

Sample Preparation and Chemical Analysis

The plants (in whole) were harvested after 5 days of inoculation. After rinsing with DI water for four times, plants were taken on clean absorbent paper to remove water from plant surfaces. The samples were then placed into a drying oven at 65 °C until they reached a constant weight. Dried samples were weighed and 0.10-0.20-g samples were digested in 50-ml polyethylene tubes (*DigiTubes*, SCP Science, Canada). Five ml of 65% HNO_3 were added and the samples were kept under a fume hood for 12 hours. Then the samples were heated to 95 °C for 2 hours on a heating block (*DigiPREP*, SCP Science, Canada). After cooling to room temperature, 3 ml of 30% hydrogen peroxide were added to the digests and the samples were heated again to 105 °C for

min and then diluted to 10 ml using DI water and stored in 15-ml polythene bottles (HDPE, NALGENE[®], Nalge Nunc International, Rochester, NY).

The concentrations of arsenic and iron were analyzed using a graphite-furnace atomic absorption spectrometer (GF-AAS, Z-8100, Hitachi, Japan). For the determination of arsenic, 5 μL of 0.05M nickel nitrate was added to a 10- μL sample as matrix modifier in the cuvette. The accuracy of the analysis was checked by the analysis of certified standard reference material 1573a tomato leaf (NIST, USA). The arsenic concentration in certified reference material was $0.112 \pm 0.004 \mu\text{g g}^{-1}$ while the measured arsenic concentration was $0.123 \pm 0.009 \mu\text{g g}^{-1}$. The concentrations detected in all samples were above the instrumental limits of detection ($\geq 0.01 \mu\text{M}$ in samples in water). Total phosphate was determined spectrophotometrically [29].

Chemical reagents used in this experiment were of analytical grade. All glass wares used were washed with detergent solution, 3 M HCl and finally with DI water for eight times before use. In each analytical batch at least two reagent blanks and three replicate samples were included.

Data Analysis

The experimental data were statistically analyzed for mean separation of different arsenic treatments according to the least significant difference (LSD) at 5% level by IRRI-STAT 4.0 for windows (developed by the Biometrics unit, IRRI, Philippines) and the Pearson correlation coefficient (r) was calculated by SPSS[®] statistical package (version 10.0 for windows).

Results and Discussions:

Uptake of Arsenic Species by *Salvinia natans* L. From Culture Solution

The arsenic uptake by water fern (*Salvinia natans* L.) at different phosphate concentrations are shown in Fig. 1. After 5 days of incubation, the water fern accumulated a maximum of $0.24 \pm 0.02 \mu\text{mol (g dry weight)}^{-1}$ of arsenic from phosphate deficient solution ($P = 0 \mu\text{M}$) and a minimum of $0.14 \pm 0.02 \mu\text{mol (g dry weight)}^{-1}$ from phosphate-rich solution ($P = 100 \mu\text{M}$) when the MS culture solutions were modified with $4.0 \mu\text{M}$ of arsenate. The results imply that arsenate uptake into the water fern was significantly higher in phosphate deficient solutions than the phosphate-rich solutions and the increase of phosphate concentration in culture solution decreases arsenate uptake. However, arsenic accumulation by the plants was highest ($0.03 \pm 0.00 \mu\text{mol g}^{-1}$ dry weight) in phosphate sufficient solution ($P = 100 \mu\text{M}$) when the initial concentrations of DMAA in growth medium was $4.0 \mu\text{M}$. This concentration of arsenic in plant tissue did not differ significantly with the concentration ($0.02 \pm 0.00 \mu\text{mol g}^{-1}$ dry weight), when the plants were grown in phosphate deficient growth medium ($P = 0 \mu\text{M}$). This might be because the DMAA uptake in the aquatic macrophyte was not affected by the initial phosphate concentrations in the solution.

Phosphate added to the growth medium plays two important roles: i) it enhances arsenate availability in the solution; and, ii) it competes with arsenate for uptake carriers in the plasmalemma due to the similar chemical behavior of arsenate and phosphate [30, 31]. The fact that arsenate and phosphate concentrations in tissues of *Salvinia natans* L. were significantly negatively correlated ($r = -0.662$, $p < 0.05$) (Table 2) suggests that the competition for uptake, indeed, occurred (Fig. 2A). Mkandawire and Dudel [18] also reported that the arsenate uptake in *Lemna gibba* L. occurs through the phosphate uptake pathway due to similar chemical behavior of arsenate and phosphate. In contrast, DMAA and phosphate concentrations in tissues of *Salvinia natans* L. did not correlate significantly ($r = -0.076$, $p > 0.05$) (Fig. 2B). This is because DMAA does

not compete with phosphate for plant uptake due to their dissimilar chemical behavior.

Effect of Arsenic Species on Phosphate Uptake by *Salvinia natans* L.

Arsenate in the culture solutions significantly ($p < 0.05$) reduced phosphate uptake in tissues of *Salvinia natans* L. However, the DMAA did not affect phosphate uptake into the plant significantly ($p > 0.05$). The Pearson correlation analysis (Table 2) revealed a significant negative relationship between arsenate and phosphate concentrations in tissues of *Salvinia natans* L. (Fig. 2A). No significant correlation was observed between DMAA and phosphate concentrations in tissues of *Salvinia natans* L. (Fig. 2B). Reduction of phosphate uptake in plants exposed to arsenate has also been reported in literatures [31, 32]. This is because the arsenate uptake occurs through the phosphate uptake pathway even replacing the phosphate from sorption site [33]. The DMAA may be accumulated in *Salvinia natans* L. through different mechanisms.

Arsenic Removal Efficiency of *Salvinia natans* L.

After 5 days of exposure to culture solutions containing different concentrations of arsenate, the *Salvinia natans* L. removed a significant amount of arsenic (Fig. 3). Regardless of phosphate concentrations in solution, between 32-65% arsenate was removed from the solution by *Salvinia natans* L. within the five days for a plant dry biomass of 0.15 g. On the other hand, DMAA removal was negligible (about 0.7-3.2%). The results indicate that removal of arsenic were increased with the increase of arsenate concentrations and decreased with the increase of phosphate concentrations in the solution. Mukherjee et al. [34] reported a 74.8% removal of arsenic by the same plant within 120 hrs of exposure when the initial source of arsenic was arsenate (As(V)).

Influence of Phosphate and Iron on Arsenic Uptake in *Salvinia natans* L.

Fig. 4 shows the correlation between arsenic and iron concentrations in *Salvinia natans* L. Arsenate was found to be significantly positively correlated ($r = 0.662$; $p < 0.05$) with iron while DMAA was independent of iron concentration ($r = 0.233$; $p > 0.05$) (Table 2). Robinson et al. [33] also found a positive correlation between arsenic and iron in native aquatic ferns (*Asplenium bulbiferum*, *Blechnum discolor*, *Histiopteris incisa*, *Pneumatopteris penningera* and *Polystichum vestitum*) as well as watercress (*Rorippa nasturium-aquaticum*). This might be due to the physico-chemical adsorption of arsenate on iron oxides on plant surfaces. Robinson et al. [33] discussed the physico-chemical as an alternative mechanism of arsenic accumulation in aquatic plants. In this mechanism, iron oxides (iron plaques) on the plant surfaces adsorb and accumulate arsenic. Although arsenic adsorption on iron oxide plaques on the surface of aquatic plants has been reported by Robinson et al. [33], which species of arsenic predominated in such adsorption was not clear from their studies. However, Blute et al. [35] reported arsenate to be positively correlated with iron plaques on roots of *Typha latifolia* (cattail) grown in arsenic-contaminated wetland sediments. According to Blute et al. [35], the ferric plaques were predominantly Fe(III) oxyhydroxide and 80% of the arsenic in it were arsenate. The present study demonstrates that arsenic adsorbed on the iron plaques of aquatic plant surfaces is mainly arsenate, as it was adsorbed on iron plaques of wetland plant *Typha latifolia* (cattail).

Arsenate and iron concentrations in *Salvinia natans* L. were highly positively correlated ($p < 0.01$) when the plants were grown in phosphate-deficient solution while their correlation was not significant ($p > 0.05$), when the plants were grown in phosphate-sufficient solution. The result suggests that phosphate is adsorbed on iron oxides (Fe-plaques) of aquatic plant surfaces and displace arsenate from the sorption sites on iron oxides. It is well established that iron (hydr)oxides are important phosphate adsorbents in soils [36-39] oxic sediments [40]. The use of Fe oxides to

adsorb phosphate on-site and reduce its concentrations in runoff and leachates is a proven approach to potentially lowering phosphate loadings of water bodies [41-43]. Numerous laboratory studies have also been directed at the sorption of phosphate on Fe oxides [44-47].

Some studies have attempted to quantify differences in phosphate adsorption associated with variations in mineral properties such as surface area, morphology, and chemical composition [47, 48]. Ferrihydrite is perhaps the most effective of these minerals in terms of phosphate adsorption in soils due to its small particle size, high surface area, and gel-like form. In nature, ferrihydrite is formed by the rapid oxidation of Fe(II) in Fe-rich waters [49]. Thus, the phosphate provably not only compete with arsenate for uptake carriers in plasmalemma [17] but also compete for adsorption on iron oxides of roots or plant surfaces as the phosphate and arsenate are analogous in chemical properties. The competition between arsenate and phosphate for the adsorption on iron oxides of plant surfaces results in the reduction of physico-chemical adsorption of arsenate in aquatic plants.

Conclusion:

Phosphate and iron are two important nutrient elements affecting the arsenic uptake in water fern *Salvinia natans* L. The *Salvinia natans* L. uptake arsenate probably through symplastic or apoplastic pathway and compete with phosphate for uptake carriers in plasmalemma. But stronger binding affinity of phosphate with the uptake carriers inhibits arsenate uptake in aquatic plants. However, physicochemical adsorption would be an alternative and potential mechanism for arsenic uptake in aquatic plants. In this mechanism, arsenate is adsorbed by iron oxides on plant surfaces. Although the present study reveals the physicochemical uptake of arsenate in water fern, the individual concentrations of arsenic in plant tissue and iron plaques were not measured. Therefore, it is

difficult to interpret how much arsenic and iron was taken up in the plant tissues. It needs microanalysis of the tissues to make the fact clear. But as iron (hydr)oxides are important phosphate adsorbents and the phosphate has stronger binding affinity to the uptake carriers in plasmalemma, low correlation coefficient between arsenate and iron in plants of phosphate-sufficient solution suggest that most of the arsenate might be bound to the outer cell wall rather than entering into the plant tissues. Nevertheless, this does not decrease the importance of aquatic macrophytes in arsenic phytoremediation.

Acknowledgements:

This research was supported partly by Grants-in-Aid for Scientific Research (18510071) from the Japan Society for the Promotion of Science, and the Steel Industry Foundation for the Advancement of Environmental Protection Technology, Japan.

References:

- 1) M. A. [Rahman](#), H. Hasegawa, K. Ueda, T. Maki, C. Okumura, M. M. Rahman, Arsenic accumulation in duckweed(*Spirodela polyrhiza* L.): A good option for phytoremediation. Chemosphere. 69 (2007) 493-499.
- 2) M. A. [Fazal](#), T. Kawachi, E. Ichio, Validity of the latest research findings on causes of groundwater arsenic contamination in Bangladesh. Water International. 26 (2001) 380-89.
- 3) A. H. [Smith](#), E.O. Lingas, M. Rahman, Contamination of drinking water by arsenic in Bangladesh: a public health emergency. Bull. Of the World Health Organization, 78(2000) 1093-1103.

- 4) K. M. [Ahmed](#), Groundwater arsenic contamination in Bangladesh: An overview. In: Bhattacharya, P. and Welch, A. H. (Eds.). Arsenic in groundwater of sedimentary aquifers. 31st International geological congress, Rio de Janeiro, Brazil. (2000) 3-11.
- 5) A. K. [Chakraborti](#), D. K. Das, Arsenic pollution and its environmental significance. J. Interacad 1 (1997) 262-276.
- 6) D. M. [Banerjee](#), Some comments on the source of arsenic in the Bengal Deltaic sediments. In: Bhattacharya, P. and Welch, A. H. (Eds.). Arsenic in groundwater of sedimentary aquifers. 31st International geological congress, Rio de Janeiro, Brazil. (2000) 15-17.
- 7) P. [Visoottiviseth](#), K. Francesconi, W. Sridokchan, The potential of Thai indigenous plant species for the phytoremediation of arsenic contaminated land. Environ. Poll. 118 (2002) 453-461.
- 8) P. [O'Neill](#), Arsenic, In: Heavy metals in soils. B. J. Alloway (Ed.). (1995) 105-121.
- 9) P. L. [Smedley](#), D. G. Kinniburgh, A review of the source, behaviour and distribution of arsenic in natural waters. Appl. Geochem. 17 (2002) 517-568.
- 10) I. [Raskin](#), P. B. A. Nanda-Kumar, S. Dushenkov, D. E. Salt, B. D. Ensley, Removal of radionuclides and heavy metals from water and soil by plants. OECD Document, Bioremediation. (1994) 345-354.
- 11) T. [De Koe](#), *Agrostis castellana* and *Agrostis delicatula* on heavy metal and arsenic enriched sites in NE Portugal. Sci. Total. Envi. 145 (1994) 103-109.
- 12) J. [Bech](#), C. Poschenrieder, M. Llugany, J. Barcelo, P. Tume, F.J. Toloias, As and heavy metal contamination of soil and vegetation around a copper mine in Northern Peru. Sci. Total Envi. 203 (1997) 83-91.

- 13) L. Q. [Ma](#), K. M. Komar, C. Tu, W. Zhang, Y. Cai, E. D. Kennelley, A fern that hyperaccumulates arsenic. *Nature*. 409 (2001) 579.
- 14) P. A. [Gulz](#), S. K. Gupta, R. Schulin, Arsenic accumulation of common plants from contaminated soils. *Plant and soil*. 272 (2005) 337-347.
- 15) K. [Komar](#), L. Q. Ma, D. Rockwood, A. Syed, Identification of arsenic tolerant and hyperaccumulating plants from arsenic contaminated soils in Florida. *Agron. Abstr.*(1998) 343.
- 16) C. [Tu](#), L. Q. Ma, B. Bondada, Arsenic accumulation in the hyperaccumulator Chinese brake and its utilization potential for phytoremediation. *J. Environ. Qual.* 31 (2002) 1671-1675.
- 17) M. [Mkandawire](#), Y. V. Lyubun, P. V. Kosterin, E. G. Dudel, Toxicity of arsenic species to *Lemna gibba* L. and the influence of phosphate on arsenic bioavailability. *Environ. Toxicol.* 19 (2004) 26-35.
- 18) M. [Mkandawire](#), E. G. Dudel, Accumulation of arsenic in *Lemna gibba* L. (duckweed) in tailing waters of two abandoned uranium mining sites in Saxony, Germany. *Sci. Total Environ.* 336 (2005) 81-89.
- 19) C. K. [Lee](#), K. S. Low, N. S. Hew, Accumulation of arsenic by aquatic plants. *Sci. Total Environ.* 103 (1991) 215-227.
- 20) B. [Robinson](#), C. Duwing, N. Bolan, M. Kannathasan, A. Saravanan, Uptake of arsenic by New Zealand watercress (*Lepidium sativum* L.). *Sci. Total Environ.* 301 (2003) 67-73.
- 21) O. I. [Sizova](#), V. V. Kochetkov, S. Z. Validov, A. M. Boronin, P. V. Kosterin, Y. V. Lyubun, Arsenic-contaminated soils: genetically modified *Pseudomonas* spp. and their arsenic-phytoremediation potential. *J. Soils and Sediments*. 2 (2002) 19-23.

- 22) P. [Selvapathy](#), P. Sreedhar, Heavy metals removal by water hyacinth. J. Indian Publ. Health Engr. 3 (1991) 11-17.
- 23) B. [Alam](#), A. K. Chatterjee, S. Duttagupta, Bioaccumulation of Cd (II) by water lettuce. Pollut. Res. 14 (1995) 59-64.
- 24) A. K. [Sen](#), N. G. Mondal, *Salvinia natans* as the scavenger of Hg (II). Wat. Air Soil Pollut. 34 (1987) 439-446.
- 25) A. K. [Sen](#), M. Bhattacharyya, Studies on uptake and toxic effects of lead on *Salvinianatans*. Indian J. Environ. Health. 35 (1993) 308-320.
- 26) K. S. [Low](#), C. K. Lee, C. H. Tai, Biosorption of copper by water hyacinth roots. J. Environ. Sci. Health A. 29(1) (1994) 171-188.
- 27) N. W. [Ingole](#), J. P. Ting, Study on nutrient removal potential of selected aquatic macrophytes. J. Inst. Engr (India) Environ. Engng Div. 83 (2002) 1-6.
- 28) A. K. [Sen](#), N. G. Mondal, Removal and uptake of copper (II) by *Salvinia natans* from wastewater. Wat. Air Soil Pollut. 49 (1990) 1-6.
- 29) S. C. [Lenore](#), E. G. Arnold, D. E. Andrew, (Eds.), Standard methods for the examination of water and wastewater, 20th ed. APHA, AWWA and WEF. 1998.
- 30) C. [Tu](#), L.Q. Ma, Effects of arsenic and phosphate on their accumulation by an arsenic-hyperaccumulator *Pteris vittata* L. Plant Soil, 249 (2003) 373-382.
- 31) J. [Wang](#), F. J. Zhao, A. A. Meharg, A. Raab, J. Feldman, S. P. McGrath, Mechanism of arsenic hyperaccumulation in *Pteris vittata* L. uptake kinetics, interaction with phosphate and arsenic speciation. Plant Physiol. 130 (2002) 1552-1561.

- 32) M. [Patra](#), N. Bhowmil, B. Bandopadhyay, A. Sharma, Comparison of mercury, lead and arsenic with respect to genotoxic effects on plant systems and the development of genetic tolerance. Environ. Exp. Bot. 52 (2004) 199-223.
- 33) B. [Robinson](#), N. Kim, M. Marchetti, C. Moni, L. Schroeter, C. van den Dijssel, G. Milne, B. Clothier, Arsenic hyperaccumulation by aquatic macrophytes in the Taupo Volcanic Zone, New Zealand. Environ Exper Bot. 58 (2006) 206-215.
- 34) S. [Mukherjee](#), S. Kumar, Adsorptive uptake of arsenic (V) from water by aquatic fern *Salvinia natans*. J. Water Supply: Research and Technology- AQUA. 54(1) (2005) 47-53.
- 35) N. K. [Blute](#), D. J. Brabander, H. F. Hemond, S. R. Sutton, M. G. Newville, M. L. Rivers, Arsenic sequestration by ferric iron plaque on cattail roots. Environ. Sci. Technol. 38(2004) 6074-6077.
- 36) G. [Guzman](#), E. Alcantara, V. Barron, J. Torrent, Phytoavailability of phosphate adsorbed on ferrihydrite, hematite, and goethite. Plant and Soil. 159 (1994) 219-225.
- 37) U. [Schwertmann](#), R. M. Taylor, Iron oxides. In: Minerals in Soil Environments (2nd ed.), B. Dixon, S. B. Weed (Eds.), Soil Sci. Soc. Am. Madison, Wisconsin, USA. (1989) 378-438.
- 38) O. K. [Borggaard](#), S. S. Jørgensen, J. P. Moberg, B. Raben-lange, Influence of organic matter on phosphate adsorption by aluminium and iron oxides in sandy soils. European J. Soil Science. 41 (1990) 443-449.
- 39) M. E. [Hamad](#), D. L. Rimmer, J. K. Syers, Effect of iron oxide on phosphate sorption by calcite and calcareous soils. European J. of Soil Science. 43 (1992) 273-281.
- 40) M. D. [Krom](#), R. A. Berner, Adsorption of phosphate in anoxic marine sediments. Limnol. Oceanogr. 25 (1980) 797-806.

- 41) P. A. [Moboornea](#), D. M. Miller, Decreased phosphorus solubility in poultry litter with aluminum, calcium, and iron amendments. *J. Environ. Qual.* 23 (1994) 325-330.
- 42) H. A. [Elliott](#), G. A. O'Connor, P. Lu, S. Brinton, Influence of water treatment residuals on phosphorus solubility and leaching. *J. Environ. Qual.* 31 (2002) 1362-1369.
- 43) L. E. [Gallimore](#), N. T. Basta, D. E. Storm, M. E. Payton, R. H. Huhnke, M. D. Smolen, Water treatment residual to reduce nutrients in surface runoff from agricultural land. *J. Environ. Qual.* 28 (1999) 1474-1478.
- 44) R. L. [Parfitt](#), Anion adsorption by soils and soil materials. *Adv. Agron.* 30 (1978) 1-50.
- 45) J. B. [Harrison](#), V.B. Berkheiser, Anion interactions with freshly prepared hydrous ironoxides. *Clays Clay Miner.* 30 (1982) 97-102.
- 46) C. C. [Ainsworth](#), M. E. Sumner, V. J. Hurst, Effect of aluminum substitution in goethite on phosphorus adsorption: I. Adsorption and isotopic exchange. *Soil Sci. Soc. Am. J.* 49 (1985) 1142-1149.
- 47) V. [Barron](#), M. Herruzo, J. Torrent, Phosphate adsorption by aluminous hematites of different shapes. *Soil Sci. Soc. Am. J.* 52 (1988) 647-651.
- 48) J. [Torrent](#), V. Barron, U. Schwertmann, Phosphate adsorption and desorption by goethites differing in crystal morphology *Soil Sci. Soc. Am. J.* 54 (1990) 1007-1012.
- 49) F. E. [Rhoton](#), J. M. Bigham, D. L. Lindbo, Properties of iron oxides in streams draining the loess uplands of Mississippi. *Appl. Geochem.* 17 (2002) 409-419.

Table 1: Modified^a Murashige and Skoog (MS) culture solution used for *Salvinia natans* L. cultivation.

Nutrient	Concentration (mg L ⁻¹)
KNO ₃	1900.00
NH ₄ NO ₃	1650.00
CaCl ₂ .2H ₂ O	440.00
MgSO ₄ .7H ₂ O	370.00
K ₂ HPO ₄	Modified ^a
FeSO ₄ .7H ₂ O	27.80
MnSO ₄ .5H ₂ O	22.30
ZnSO ₄ .7H ₂ O	8.60
H ₃ BO ₃	6.20
KI	0.83
Na ₂ MoO ₄ .2H ₂ O	0.25
CuSO ₄ .5H ₂ O	0.025
CoCl ₂ .6H ₂ O	0.025
Na ₂ -EDTA	37.30

^a The PO₄³⁻ concentration in control treatment was 0.02 µM and the other concentrations of PO₄³⁻ in the solutions were 100 and 500 µM.

Table 2: Correlations between arsenic, phosphate, and iron uptake in *Salvinia natans* L. exposed to arsenate and DMAA.

Correlation between	Pearson Correlation (r)	Significance (<i>p</i>)
As(V) & P	-0.662*	0.019
DMAA & P	-0.076	0.814
As(V) & Fe	0.662*	0.019
DMAA & Fe	0.233	0.466

* Correlation is significant at the 0.05 level




Figure 1: Arsenic uptake in *Salvinia natans* L. affected by the phosphate concentrations in culture solution.

Error bars represent \pm S.D. ($n = 3$).




Fig. 2: Rerelationship between arsenic and phosphate concentrations in of *Salvinia natans* L. exposed to arsenate (A) and DMAA (B).




Fig. 3: Influence of phosphate on arsenic uptake in *Salvinia natans* L. exposed to different concentrations of DMAA. Error bars represent \pm S.D. ($n = 3$).




Fig. 4: Correlation between arsenic and iron concentrations in of *Salvinia natans* L. exposed to arsenate (A) and DMAA (B).