

Examination of factors affecting gait properties in healthy older adults: Focusing on knee extension strength, visual acuity, and knee joint pain

メタデータ	言語: eng 出版者: 公開日: 2017-10-02 キーワード (Ja): キーワード (En): 作成者: メールアドレス: 所属:
URL	http://hdl.handle.net/2297/39047

Original Article

Title:

**Examination of factors affecting gait properties in healthy elderly people
- focusing on knee extension strength, visual acuity, and knee joint pain –**

Tomohiro Demura¹, Shin-ichi Demura², Masanobu Uchiyama³, Hiroki Sugiura²

¹ *Child Education, Jin-ai Women's College,*

Fukui, Fukui, Japan

² *Graduate School of Natural Science & Technology, Kanazawa University,*

Kanazawa, Ishikawa, Japan

³ *Akita Prefectural University,*

Shimoshinjo-Nakano, Akita, Japan

***Corresponding author:** Tomohiro Demura

Jin-ai Women's College

Child Education,

Amaiike, Fukui, Fukui 910-0134, Japan

Tel : +81-0776-56-1133, Fax: +81-0776-56-2922

E-mail: tomodemu@jin-ai.ac.jp

Background & Purpose: Gait properties change with age due to a decrease in leg strength and visual acuity or knee joint disorders. Gait changes commonly result from these combined factors. This study aimed to examine the effects of knee extension strength, visual acuity, and knee joint pain on gait change for 181 healthy female *older adults* (age: 76.1 +/- 5.7 years).

Methods: Walking speed, cadence, stance time, swing time, double support time, step length, step width, walking angle and toe angle were selected as gait parameters. Knee extension strength was measured by isometric knee extension strength, and decreased visual acuity and knee joint pain were evaluated by subjective judgment whether or not such factors created a hindrance during walking.

Results: Among *older adults* without vision problems and knee joint pain that affected walking, those with superior knee extension strength had significantly greater values of walking speed and step length than those with inferior knee extension strength ($p < 0.05$).

Persons with visual acuity problem had more cadence and shorter stance time. In addition, persons with pain in both knees showed slower walking speed and longer stance time and double support time.

Conclusion: A decrease of knee extension strength and visual acuity, and knee joint pain are factors affecting gait in the female *older adults*.

Decreased knee extension strength and knee joint pain mainly affect respective distance and time parameters of the gait.

Keywords: Gait; Human locomotion; *Older adults*

Introduction

Walking is an important element for various daily living activities. Gait properties that can assess walking quantitatively may be a simple measure for evaluating overall human health.¹ In old age, various physical functions including leg strength, joints, and visual acuity, decrease and greatly affect gait. It was reported that a gait change with age also relates to falls.²⁻⁴ In addition, gait has been used as an indicator for predicting the risk of disease and death in *older adults*.³

As concrete parameters of gait properties, generally, stance time, swing time, step length, and step width including walking speed and cadence, have been used.^{5,6} The following changes have been reported to occur in walking with aging: a decrease in walking speed,^{5,7} an increase in single leg support time and a decrease in double leg support time during total contact time,⁸ and shortness of step length.⁹

The main factors related to gait change in old age include leg muscle function,¹⁰⁻¹² vision function,^{13,14} and knee joint function.^{15,16} Among others, there are many reports concerning the effect of a decrease in leg strength on gait.
10-12

Visual information is important for stable walking, and a decrease of visual function (e. g. low visual acuity) produces disadvantage in achievement of daily living activities.¹⁷ In the case of *older adults*, it was reported that a decrease in visual acuity increases the risk of falls.¹⁸⁻²⁰ Elliott et al.²¹ confirmed that walking speed and cadence decreased when young healthy persons with

simulated cataracts walked in dim light, and Moe-Nilssen et al.²² reported that a sudden reduction from normal to marginal lighting while walking induced gait change.

Additionally, leg joints play an important role in gait as walking is a movement that primarily requires use of the lower limbs. In old age, knee joint function decreases, and patients with knee osteoarthritis have slower walking speed, and slower cadence, or shorter step length, and longer stance time.^{23,24} Kaufman et al.²⁵ confirmed that compared with healthy subjects, knee osteoarthritis patients walk with a lower knee extension moment to reduce knee strain. Generally, *older adults* have suppressed joint function. About 25% of *older adults* aged 55 or over, including knee osteoarthritis patients, complain of knee joint pain.^{26,27} Knee joint pain greatly affects walking.²⁸

In this way, it is considered that vision problems and knee joint pain, in addition to decreased leg strength are important factors affecting gait.

In old age, it is commonly the case that many physical functions decrease simultaneously. In short, *older adults* generally have some problems (e. g. decrease of leg strength and vision function, and knee joint pain) which affect gait. When examining *older adults* with vision problems and knee joint pain in addition to leg strength as subjects, it is difficult to identify the specific factor that affects a gait change. Not only leg strength, but also vision problem or knee joint pain may be involved in gait change.

This study aimed to examine the effects of knee extension strength, visual acuity, and knee joint pain on gait change in *older adults* by considering synergies among these factors.

Materials and Methods

Subjects

The subjects that participated in this study were 181 healthy, community-dwelling female *older adults*. Table 1 shows their age and physical characteristics. All subjects participated in health class or social educational activities that were hosted by municipal governments. In addition, their ADL score was very high (Ministry of Education, Culture, Sports, Science & Technology). Hence, they were judged to have high independence in daily activities because many of them engaged in leisure activity and work. Before the measurements, the purpose and procedure of this study were explained in detail and informed consent was obtained from all subjects. In addition, this study was approved by the Ethics Committee on Human Experimentation of Faculty of Education, Kanazawa University (Ref. No. 19-17).

*** Table 1 near here ***

Testing Protocol

Gait Property

Gait properties were measured by a gait analysis system Walk Way MG-1000 (Anima, Japan) referenced to a Demura and Demura's ⁶ method. The MG-1000 with plate sensors can determine time, dimensions and the distance of the foot or feet when the foot touched its surface and can measure

grounding/non-grounding on the bearing surface as an on/off signal. Data were recorded into a personal computer at 100 Hz. Subjects walked 12m road (wooden floor) with above sensors with voluntary maximum speed. The analysis interval was from 4m to 8m.

Factors

Leg strength:(The effect of knee extension strength

Knee extension strength was selected for assessing leg strength.²⁹ During measurement of isometric knee extension strength, the subjects were seated upright in a rigid chair with knees flexed an angle of 90° and with the lower legs strapped in a pad just above the ankle, attached by a backward rigid bar to a tension meter attachment (T.K.K.1269f; Takei Scientific Instruments Co. Ltd., Japan). In addition, the subjects folded their arms on their chest. The subject was asked to extend the knee as far as possible and to maintain it for 3 seconds. Each of the subjects' legs was measured twice at intervals of greater than thirty seconds. It was difficult to adequately judge whether knee extension strength of *older adults* is superior or inferior because there were little data on their knee extension strength measured with the same method. Hence, superior and inferior groups in knee extension strength were made up among subjects measured in this study. In short, based on mean value (Mean) and standard deviation (SD), subjects were divided into three groups with different knee extension strength (Inferior group: Inferior < Mean-1SD, Middle group: Mean-1SD < Middle < Mean+1SD, Superior group: Mean+1SD < Superior)

Visual acuity: The effect of subjective visual acuity problems

In old age, a decrease in visual function greatly affects gait. Persons who were selected in this study for visual acuity found it difficult to read characters or see people's faces, and felt hindered in their gait. Many of **older adults** interfere with walking due to narrow visual field even if having normal eyesight. Hence, in this study, it was judged that there is an adequate method to ask whether **older adults** feel totally trouble in actual walking. Incidentally, such index about visual acuity has been used also in fall risk survey.³⁰

Knee joint pain: The effect of subjective knee joint pain

Having knee joints with orthopedic disorders affect walking.^{23,24} It was also assumed that subjective knee pain affects walking.²⁸ Referring Sugiura and Demura's¹ method, Persons with knee joint pain were divided into those who had pain in one knee joint or those who had pain in both knee joints.

Parameters

Gait variability (Temporal and spatial parameters)

The following 9 gait parameters were selected in reference to previous studies:⁶ walking speed, cadence, stance time, swing time, double support time, step length, step width, walking angle, and toe angle.

Walking speed: walking distance per second (cm/sec).

Cadence: the number of steps per minute (steps/min).

Stance time: the duration that the body is supported by one or both feet, that is, the phase in which one foot or both feet contact the floor.

Swing time: the duration that one foot swings, that is, one foot leaves the floor. This time agrees with a single support time (sec).

Double support time: the duration in which both feet contact the floor (sec).

Step length: the distance between anterior-posterior patterns (one step length).

Step width: the distance between both feet (cm).

Walking angle: the angle between the direction of movement and bilateral pattern line ($^{\circ}$).

Toe angle: the angle between the direction of movement and the foot axis ($^{\circ}$).

Data Analysis

Older adults usually have a decrease in knee extension strength and vision problems or else knee joint pain simultaneously. The effect of the other factors was controlled to examine the unique effect of each factor.

The effect of knee extension strength on gait (F1)

To examine effect of leg strength on gait, mean differences of gait parameters among three groups with different knee extension strength (inferior, middle, and superior, divided by mean and SD) for subjects without a visual acuity problem and knee joint pain were tested using one-way analysis of variance (ANOVA).

The effect of subjective visual acuity problem on gait (F2)

To examine effect of the visual acuity problem on gait, mean differences in gait parameters between groups with/without visual acuity problems for subjects without knee joint pain were tested using a t-test.

The effect of subjective knee joint pain on gait (F3)

To examine effect of knee joint pain on gait, mean differences of gait parameters between groups with/without knee joint pain for subjects without visual acuity problem were tested using one-way analysis of variance (ANOVA).

In addition, if significant differences were found in age, height, weight, and knee extension strength, gait parameters were tested using analysis of covariance. A Tukey's HSD (Honestly Significant Difference) method was selected for multiple comparisons. The probability level of 0.05 was indicative of statistical significance.

Results

The effect of knee extension strength on gait (F1)

Table 2 shows the basic statistics and results of one-way analysis of variance (ANOVA) for gait parameters among three groups with different knee extension strength (inferior, middle, and superior, divided by mean and SD) for subjects without visual acuity problem and knee joint pain. Walking speed and

step length was significantly larger in the superior group than in the middle and inferior groups.

*** Table 2 near here ***

The effect of subjective visual acuity problems on gait (F2)

Table 3 shows the basic statistics and results of the t-test for gait parameters between those with and without visual acuity problems for subjects without knee joint pain. Persons with visual acuity problems as compared with those without them showed a larger value in cadence and a less value in stance time.

*** Table 3 near here ***

The effect of subjective knee joint pain on gait (F3)

Table 4 shows the basic statistics and results of one-way ANOVA for gait parameters among the three groups with different knee joint pain (one knee, both knee, and none) for subjects without visual acuity problems. Because significant differences were found in body mass and knee extension strength analysis of covariance was used. Walking speed is faster in persons with pain in both knees as compared with persons without knee pain. Persons with pain in both knees as compared with persons with pain in one knee or no pain showed a larger value in stance time and double support time.

*** Table 4 near here ***

Discussion

The effect of knee extension strength on gait

In examining for **older women** without visual acuity problems and knee joint pain, those with superior knee extension strength had significantly larger values in walking speed and step length than those with inferior knee extension strength. It has been reported that gait properties in **older adults** include a decrease in step length and an increase in double support time and step width.^{32,33} Patla³⁴ reported that these gait properties in **older adults**, which differ from younger people, may contribute to increased walking stability.

A decrease in leg strength is closely related to a decrease in walking speed and is considered to be a main cause of gait change with age.^{35,36} Ferrandez et al.³⁷ reported that if walking speed is the same, gait movements in **older adults** will not be largely different from those of young adults. From the above, similarly to results of previous studies,¹⁰⁻¹² it is judged that knee extension strength contributes closely to a decrease in walking speed and step length.

The effect of subjective visual acuity problem on gait

It was reported that fall risk in **older adults** increases with a decrease in visual acuity.^{18,38,39} However, there are also reports that there was no relationship between visual acuity and the fall risk when considering factors such as age.^{3,40}

On the other hand, walking speed decreased when young persons with simulated cataract walked in dim light.²¹ Furthermore, a sudden change from normal to marginal lighting induces gait change, but gait characteristics are nearly normalized due to the adaptation to darkness during the first 90 seconds, suggesting that the darkness itself does not cause the gait changes.²² Also in this study, cadence was more and stance time was shorter due to visual acuity problem. However, the above will be necessary to examine carefully because *older women* with visual acuity problems as compared with *older women* without them showed faster walking speed and wider step length, although significance was not found. In addition, the following may affect also the present result; there was no task in which subjects recognize an obstacle in which visual acuity is affected due to using a flat walkway in this study.

The effect of subjective knee joint pain on gait

Al-Zahrani and Bakheit²³ and Bejek et al.²⁴ reported that osteoarthritis patients show a decrease in walking speed and step length, and an increase in stance time. In addition, an increase in step width is found as one of the features of abnormal gait in people with disorders.³³ Gehlsen and Whaley⁴¹ reported that *older adults* with fall history have significantly larger step width than those without it. Hence, step width may be an efficient means of judging abnormal gait.

Moreover, Bejek et al.²⁴ reported that osteoarthritis patients, when walking with fast speed compared to slow speed, show a more prominent gait change due

to the effects of pain.

In this study, *older women* with pain in both knees showed slow walking speed and long stance time and double support time. This gait changes are similar to those reported by Bejek et al.²⁴ In addition, it is considered to be one of characteristics in persons with subjective knees joint pain because the above parameters are mainly ones regarding time.

Conclusion

This study examined the effects of knee extension strength, subjective visual acuity and subjective knee joint pain on gait change in *older adults*.

Older adults with superior knee extension strength can walk at a faster speed and with longer step length than *older adults* with inferior knee extension strength. Persons with visual acuity problems had more cadence and shorter stance time. In addition, persons with pain in both knees showed slower walking speed and longer stance time and double support time.

In conclusion, a decrease of knee extension strength and visual acuity, and knee joint pain are factors affecting gait in the female *older adults*. Decreased knee extension strength and knee joint pain mainly affect respective distance and time parameters of the gait.

Acknowledgements

Research funds were not provided by any institution.

References

1. Tang hai-peng, Toyoshima S, Hoshikawa T. Characteristics in walking kinematics of elderly before one year of death. *J Soc Biomech Jpn.* 2002; 26(1): 40-44. [In Japanese with English Abstract]
2. Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallage RB. Lower-extremity function in persons over the age of 70 years as a predictor of subsequent disability. *N Eng J Med.* 1995; 332: 556-561.
3. Gerdhem P, Ringsberg KA, Akesson K, Obrant KJ. Clinical history and biologic age predicted falls better than objective functional tests. *J Clin Epidemiol.* 2005; 58: 226-232.
4. Ganz DA, Bao Y, Shekelle PG, Rubenstein LZ. Will my patient fall? *JAMA.* 2007; 297(1): 77-86.
5. Murray MP, Kory RC, Clarkson BH. Walking patterns in healthy old men. *J Gerontol.* 1969; 24: 169-178.
6. Demura T, Demura S. Relationship among gait parameters while walking with varying loads. *J Physiol Anthropol.* 2010; 29 (1), 29-34.
7. Murray MP, Kory RC, Sepic SB. Walking patterns of normal women. *Arch Phys Med Rehabil.* 1970; 51: 637-650
8. Kaneko M, Morimoto Y, Kimura M, Fuchimoto K, Fuchimoto T. A kinematic analysis of walking and physical fitness in elderly women. *Can J Sport Sci.* 1991; 16(3): 223-228
9. Larish DD, Martin PE, Mungiole M. Characteristic patterns of gait in the

- healthy old. *Ann N Y Acad Sci.* 1988; 515: 18-32.
10. Murray MP, Duthie EH, Gambert SR, Sepic SB, Mollinger LA. Age-related differences in knee muscle strength in normal women. *J Gerontol.* 1985; 40(3): 275-280.
 11. Frontera WR, Hughes VA, Lutz KJ, Evans WJ. A cross-sectional study of muscle strength and mass in 45- to 78-yr-old men and women. *J Appl Physiol.* 1991; 71(2): 644-650.
 12. Manini TM, Clark BC, Nalls MA, Goodpaster BH, Ploutz-Snyder LL, Harris TB. Reduced physical activity increases intermuscular adipose tissue in healthy young adults. *Am J Clin Nutr.* 2007; 85(2): 377-384.
 13. Marron JA, Bailey IL. Visual factors and orientation-mobility performance. *Am J Optom Physiol Opt.* 1982; 59(5): 413-426.
 14. Coleman AL, Stone K, Ewing SK, Nevitt M, Cummings S, Cauley JA, Ensrud KE, Harris EL, Hochberg MC, Manqione CM. Higher risk of multiple falls among elderly women who lose visual acuity. *Ophthalmology.* 2004; 111(5): 857-862.
 15. Andriacchi TP, Andersson GB, Fermier RW, Stem D, Galante JO. A study of lower-limb mechanics during stair-climbing. *J Bone Joint Surg Am.* 1980; 62: 749-757.
 16. Riener R, Rabuffeti M, Frigo C. Stair ascent and descent at different inclinations. *Gait Posture.* 2002; 12: 143-155.
 17. Singh MM, Malhotra HS. Falls in the elderly – clinician’s approach. *J Indian Med Assoc.* 2003; 101(7): 420, 422, 424 passim.

18. Nevitt M, Comings S, Kidd S, Black D. Risk factors for recurrent non-syncopal falls. *JAMA*. 1989; 261: 2663-2668.
19. Lord SR, Ward JA, Williams P, Anstey K. Physiological factors associated with falls in older community-dwelling women. *J Am Geriatr Soc*. 1994; 42: 1110-1117.
20. Klein BEK, Moss SE, Klein R, Lee KE, Cruikshanks KJ. Associations of visual function with physical outcomes and limitations 5 years later in an older population. *The Beaver Dam Eye Study*. 2003; 110: 644-650.
21. Elliott DB, Bullimore MA, Patla AE, Whitaker D. Effect of a cataract simulation on clinical and real world vision. *Br J Ophthalmol*. 1996; 80(9): 799-804.
22. Moe-Nilssen R, Helbostad JL, Akra T, Birdedal L, Nygaard HA. Modulation of gait during visual adaptation to dark. *J Mot Behav*. 2006; 38: 118-125.
23. Al-Zaharni KS, Bakheit AM. A study of the gait characteristics of patients with chronic osteoarthritis of the knee. *Disabil Rehabil*. 2002; 24 (5): 275-280.
24. Bejek Z, Paroczai R, Illyes A, Kiss RM. The influence of walking speed on gait parameters in healthy people and in patients with osteoarthritis. *Knee Surg Sports Traumatol Arthrosc*. 2006; 14(7): 612-622.
25. Kaufman KR, Hughes C, Morrey BF, Morrey M, An KN. Gait characteristics of patients with knee osteoarthritis. *J Biomech*. 2001; 34(7): 907-915.
26. McAlindon TE, Cooper C, Kirwan JR, Dieppe PA. Knee pain and disability in the community. *Br J Rheumatol*. 1992; 31: 189-192.

27. Peat G, McCarney P, Croft P. Knee pain and osteoarthritis in older adults: a review of community burden and current use of primary health care. *Ann Rheum Dis.* 2001; 60: 91-97.
28. Dieppe P, Lim K. Osteoarthritis and related disorders: clinical features and diagnostic problems. In: Dlippel JH, Kieppe PA, eds. *Rheumatology*. 2nd ed. Ipswich (UK): Mosby; 1998: 8.3.1-10.5.
29. Nishijima T, Koyama R, Naito I, Hatakeyama S, Yamasaki H, Oku T. Relationship between knee extension muscle strength and walking ability in the elderly patients. *Rigakuryoho Kagaku*. 2004; 19(2): 95-99. [In Japanese]
30. Demura S, Susumu S, Yokoya T, Sato T. Examination of useful items for the assessment of fall risk in the community-dwelling elderly Japanese population. *Environ Health Prev Med.* 2010; 15: 169-179.
31. Sugiura H, Demura S. Effect of knee joint pain and knee joint disorder on knee extension strength and walking ability in the female elderly. *Adv Phys Edu.* 2012; 2(4): 139-143.
32. Hageman PA, Blanke DJ. Comparison of gait of young women and elderly women. *Phys Ther.* 1986; 66: 1382-1387.
33. Yamagishi T, Tokuda T. Geriatric gait – photometric analysis. *Jap J Reh Med.* 1975; 12: 97-104. [In Japanese with English Abstract]
34. Patla AE. Understanding the roles of vision in the control of human locomotion. *Gait Posture.* 1997; 5(1): 54-69.
35. Daubney ME, Culham EG. Lower-extremity muscle force and balance performance in adults aged 65 years and older. *Phys Ther.* 1999; 79(12):

1177-1185

36. den Ouden MEM, Schuurmans MJ, Arts IEMA, van der Schouw YT. Physical performance characteristics related to disability in older persons: A systematic review. *Maturitas*. 2011; 69(3): 208-219.
37. Ferrandez AM, Paihous J, Durup M. Slowness in elderly gait. *Exp Aging Res*. 1990; 16(2): 79-89
38. Bergland A, Jarnlo GB, Laake K. Predictors of falls in the elderly by location. *Aging Clin Exp Res*. 2003; 15: 43-50.
39. Klein BEK, Moss SE, Klein R, Lee KE, Cruikshanks KJ. Associations of visual function with physical outcomes and limitations 5 years later in an older population. *The Beaver Dam Eye Study*. 2003; 110: 644-650.
40. Chu LW, Chi I, Chiu AY. Incidence and predictors of falls in the Chinese elderly. *Ann Acad Med Singapore*. 2005; 34: 60-72.
41. Gehlsen GM, Whaley MH. Falls in the elderly: part I , gait. *Arch Phys Med Rehabil*. 1990; 71(10): 735-738.

Table 1. Characteristics of subjects (Female: 181)

n = 181		Mean	SD	Max	Min
Age	(years)	76.1	5.70	93.0	62.0
Height	(cm)	147.2	5.68	161.4	129.0
Weight	(kg)	49.8	7.90	70.9	31.0
Knee extension strength	(kg)	6.8	2.00	13.5	1.9
Prevalence of visual acuity problems	(%)	<u>With</u> : 38 (21%)		<u>Without</u> : 143 (79%)	
Prevalence of knee pain	(%)	<u>One</u> : 62 (34%), <u>Both</u> : 37 (20%), <u>None</u> : 82 (45%)			

With/Without: Persons with/without visual acuity problem,

One/Both/None: Persons with one-/both-/no knee pain

Table 2. The basic statistics and results of ANOVA for gait parameters among three groups with different knee extension strength groups

Knee extension strength (Mean +/- 1SD)		Inf (n=8)		Mid (n=46)		Sup (n=8)		ANOVA		Post Hoc Tukey's HSD	
		Mean	SD	Mean	SD	Mean	SD	F	p		
Knee ext strength	(kg)	4.21	0.48	7.06	1.13	11.0	1.58	77.5	0.00	*	Inf < Mid < Sup
Age	(years)	80.1	6.49	75.1	6.66	77.2	6.72	2.30	0.11		
Height	(cm)	145.0	4.98	146.3	6.21	147.6	3.37	0.45	0.64		
Weight	(kg)	48.6	7.92	46.7	7.34	48.9	8.06	0.47	0.63		
Walking speed	(cm/sec)	160.2	35.66	165.7	31.2	202.7	24.80	5.85	0.01	*	Inf, Mid < Sup
Cadence	(steps/min)	167.3	19.36	160.5	19.2	173.5	19.82	1.95	0.15		
Stance time		0.42	0.06	0.43	0.06	0.39	0.05	2.04	0.14		
Swing time	(sec)	0.31	0.03	0.33	0.03	0.32	0.03	1.61	0.21		
Double support time		0.05	0.02	0.06	0.02	0.04	0.01	2.35	0.10		
Step length	(cm)	57.3	9.22	61.9	9.23	70.4	7.54	5.04	0.01	*	Inf, Mid < Sup
Step width		7.97	2.76	7.02	2.76	7.26	2.09	0.48	0.62		
Walking angle	(°)	8.18	2.94	6.70	3.08	5.96	1.90	1.40	0.26		
Toe angle		2.82	4.42	3.09	3.09	4.19	3.17	0.49	0.62		

Low: < Mean-1SD, Mid: Mean-1SD < Mid < Mean+1SD, High: > Mean+1SD,

Knee ext strength: Knee extension strength, *: p < 0.05

Table 3. The basic statistics and results of t-test for gait parameters between groups with and without vision acuity problems

Can you see well? (newspaper, people's faces, etc.)	Yes (n=19)		No (n=64)		t-test	
	Mean	SD	Mean	SD	t	p
Age (years)	77.6	4.73	76.1	6.78	0.92	0.36
Height (cm)	146.3	6.40	146.3	5.71	-0.03	0.97
Weight (kg)	47.4	8.62	47.2	7.44	0.06	0.95
Knee extension strength (kg)	6.8	1.72	7.2	2.11	-0.72	0.48
Walking speed (cm/sec)	155.2	30.55	170.1	33.39	-1.74	0.09
Cadence (steps/min)	152.8	17.61	163.3	19.60	-2.10	0.04 *
Stance time	0.44	0.07	0.42	0.06	2.41	0.02 *
Swing time (sec)	0.34	0.03	0.33	0.03	1.09	0.28
Double support time	0.07	0.03	0.05	0.02	2.10	0.04
Step length (cm)	60.6	8.24	62.4	9.59	-0.75	0.46
Step width	8.07	2.83	7.19	2.66	1.25	0.22
Walking angle (°)	7.78	3.09	6.80	2.96	1.25	0.22
Toe angle	2.89	4.33	3.21	3.28	-0.35	0.73

*: $p < 0.05$

Table 4. The basic statistics and results of ANOVA for gait parameters among the three groups with knee joint pain

Knee joint pain		One (n=48)		Both (n=31)		None (n=64)		ANOVA		ANCOVA		Post Hoc
		Mean	SD	Mean	SD	Mean	SD	F	p	F	p	Tukey's HSD
Age	(years)	75.6	4.75	77.0	5.85	76.1	5.93	0.55	0.58			
Height	(cm)	148.8	5.52	147.1	6.09	146.3	5.70	2.77	0.07			
Weight	(kg)	52.2	7.88	51.8	6.90	46.5	9.45	7.73	0.00 *	covariate		
Knee ext strength	(kg)	6.8	2.04	6.2	2.14	7.2	2.11	2.36	0.10			
Walking speed	(cm/sec)	166.7	28.4	151.8	38.6	170.1	33.4	3.32	0.04 *	3.50	0.03 *	Both < None
Cadence	(steps/min)	163.3	24.7	152.9	24.3	163.3	22.7	2.58	0.08	2.45	0.09	
Stance time		0.43	0.06	0.48	0.09	0.42	0.06	5.69	0.00 *	5.26	0.01 *	Both > None, One
Swing time	(sec)	0.32	0.04	0.34	0.46	0.33	0.03	1.21	0.30	1.28	0.28	
Double support time		0.06	0.02	0.07	0.04	0.05	0.02	6.36	0.00 *	5.24	0.01 *	Both > None, One
Step length	(cm)	61.5	7.77	59.1	9.72	62.4	9.59	1.43	0.24	1.86	0.16	
Step width		7.02	2.62	8.14	2.82	7.19	2.66	1.80	0.17	1.70	0.19	
Walking angle	(°)	6.92	2.79	8.23	3.60	6.80	2.96	2.50	0.09	2.41	0.09	
Toe angle		2.67	4.02	3.55	4.02	3.21	3.28	0.58	0.56	0.74	0.48	

One: Persons with one knee pain on knee joints, Both: Persons with both knee pain on knee joints,

None: Persons without knee pain on knee joints, Knee ext strength: Knee extension strength